

AMERICAN MUSEUM *Novitates*

PUBLISHED BY THE AMERICAN MUSEUM OF NATURAL HISTORY
CENTRAL PARK WEST AT 79TH STREET, NEW YORK, NY 10024

Number 3476, 33 pp., 1 table

May 11, 2005

Family-Group Names for Bees (Hymenoptera: Apoidea)

MICHAEL S. ENGEL¹

CONTENTS

Abstract	2
Introduction	2
A Name for the Bees	3
Format	4
Family-Group Names	4
Systematics	12
Scraptrinae Ascher and Engel, new subfamily	13
Neffapini Ascher, new tribe	14
Afrodasy podini Engel, new tribe	15
<i>Afrodasy poda</i> Engel, new genus	16
Hesperapina Ascher and Engel, new subtribe	16
Macrogaleina Engel, new subtribe	16
Ancyloscelidina Engel and Michener, new subtribe	17
Acknowledgments	17
References	17
Appendix: Hierarchical Outline of Apoid Classification	28

¹ Division of Invertebrate Zoology, American Museum of Natural History; Division of Entomology, Natural History Museum and Biodiversity Research Center, and Department of Ecology and Evolutionary Biology, 1460 Jayhawk Boulevard, Snow Hall, University of Kansas, Lawrence, Kansas 66045-7523 (msengel@ku.edu).

ABSTRACT

The 173 family-group names for bees (Apoidea: Anthophila) are cataloged in chronological order. For each name the correct author, date, type genus, and combining stem are provided. The following names are considered *nomina nuda*: Phenacolletini, Ctenioschelini, †Chalicodomopsini, Liphanthini, Austropanurgini, and Hoplitini. The authorship of three names (Tapinotaspidini, Hexepeolini, and Ancyloscelidina) is corrected as each was a *nomen nudum* when first proposed, but has been subsequently made available by other authors. The following new names are proposed herein: **Scraptrinae** Ascher and Engel, new subfamily (Colletidae); **Nef-fapini** Ascher, new tribe (Andrenidae: Panurginae); **Afrodasyopodini** Engel, new tribe (Melittidae: Dasypodainae); **Afrodasyopoda** Engel, new genus; **Hesperapina** Ascher and Engel, new subtribe (Melittidae: Dasypodainae); **Macrogaleina** Engel, new subtribe (Apidae: Xylocopinae); and **Ancyloscelidina** Engel and Michener, new subtribe (Apidae: Apinae). A hierarchical outline of Apoidea classification (inclusive of the digger wasps), indicating the suggested current usage of all available family-group names is appended. The name Anthophila, as proposed by Latreille, is adopted for the bees as a whole.

INTRODUCTION

Increased phylogenetic scrutiny on the hierarchical structure of life has resulted in numerous reclassifications of organisms. Such alterations to the classification of life at all levels require the application of names for these groupings. Often such alterations result in relatively minor changes to reflect the increased understanding of organismal affinities. Whether minor or major, the application of names to these groups must follow sets of procedural rules so as to minimize nomenclatural instability, allowing the maximal retrieval of information from the vast biological literature, particularly older references with valuable data but obviously employing classificatory systems reflective of their era. Zoological nomenclature strictly regulates names in only three classes: the species-group (species and subspecies), the genus-group (genera and subgenera), and the family-group (names from superfamily to infratribe) (table 1). Many authors have attempted to catalog the correct application of names in the genus-group and every revision or monograph concerns itself with those of the species-group. Relatively little attention is paid, however, to names in the family-group and, as a result, tremendous confusion abounds about the history of such names, their correct form and authorship, and their accurate application in more modern systems.

The bees (Apoidea: Anthophila) are among the most intensely studied of aculeate Hymenoptera and have, perhaps, one of the more stable higher nomenclatures. This is in

TABLE 1
Categorical Ranks in the Family-Group

Hyperfamily (-oides) ^a
Superfamily (-oidea)
Epifamily (-oidae) ^a
Family (-idae)
Subfamily (-inae)
Supertribe (-iti) ^b
Tribe (-ini)
Subtribe (-ina)
Infratribe (-ita) ^a
Clan (-itoi) ^{a,c}
Subclan (-iloi) ^{a,c}
Infraclan (-isoi) ^{a,c}
Chaste (-isoa) ^{a,c}

Note: Ranks and suffixes are modified from those proposed by Bour and Dubois (1984) and McKenna and Bell (1997); ranks in boldface employed herein.

^a Rank not employed herein, but could be used to enrich the hierarchical classification at a later date (I cannot at present envision any utility, however, for recognizing family-group ranks below infratribe in the Apoidea).

^b Bour and Dubois (1984) suggested the suffix -inoi for the rank of supertribe; however, I have used the more commonly employed suffix -iti.

^c The ranks can only be loosely classified in the family group and names proposed for these ranks should perhaps not be considered coordinate with those of the higher, more traditional family-group categories.

large part due to the number of excellent monographic and phylogenetic treatments of generic and suprageneric taxa in the group (e.g., Michener, 1944, 1965, 1981; Ruz, 1987; Roig-Alsina and Michener, 1993; Alexander and Michener, 1995; Ascher, 2004).

The family-group names of bees were the subject of scrutiny by Michener (1986a); however, nearly 40 additional names have been proposed in the intervening years, several overlooked, older names have been discovered, and subsequent corrections have been identified, necessitating the current treatment.

The bees are included in the superfamily Apoidea along with a grade of wasps, the so-called “Sphecidae”, spheciform wasps, “digger” wasps, or more aptly apoid wasps. Family-group names for the apoid wasps were recently cataloged by Menke (1997), with additions and emendations by Antropov (2000) and Menke and Pulawski (2002). Although members of the Apoidea, these names are not cataloged herein for two reasons: (1) they have recently received the aforementioned taxonomic treatment and (2) they are, simply put, not bees. However, the classification of apoid wasps has recently been altered (e.g., Ohl, 1996; Prentice, 1998; Melo, 1999). Therefore, these names are included in the hierarchical outline of the superfamily (*vide* appendix 1) for the sake of completeness and to reflect the current usage of family-group names (and their synonyms) in a revised, multi-family system of Apoidea.

Numerous names have been proposed for bees or to denote suprageneric groups of bees, but are not based on an available genus-group name. Examples of such names include: Platiglossata Latreille, Nemoglossata Latreille, Mellifera Latreille (this name was even emended at one time to have a family-group suffix, as Melliferidae), Pepalephora Billberg, Archiapidae Friese, Proapidae Friese, Pygidalia Robertson, Apygidalia Robertson, Corbiculata Engel, Melissomorpha Engel, Paramelitturgini Patiny, etc. None of these, or other such names, are available in zoological nomenclature (ICZN, 1999a: Art. 11.7.1.1) and are, therefore, not considered herein. The various divisional names proposed by Rayment (1935)—i.e., Colletiformes (p. 16), Andreniformes (p. 226), Megachiliformes (p. 394), Xylcopiformes (p. 452), and Apiformes (p. 490)—could be considered family-group names since they are based on available genus-group names. Fortunately, family-group names based on all of these genera had previously been proposed

and thus Rayment’s names can be considered merely superfluous emendations (i.e., as *nomina vana*).

Instead, below I have considered only those names based on a genus-group name, regardless of the status of the generic name (i.e., whether or not the generic name has proved to be preoccupied, unavailable, invalid, etc.). This has resulted in a compilation of 173 family-group names for bees.

A NAME FOR THE BEES

Since the rightful unification of the apoid wasps and the bees into a single superfamily, many melittologists have lamented that the bees, themselves, now lack a formal rank. This desire to have the bees singled out with an official rank and name has led some to demote them all to a single family, Apidae. Given the diversity of bees and the tremendous utility of a multi-familial classification for this lineage, I believe it is worthwhile to retain the current classification rather than demote each taxon. However, I disagree that the bees as a whole require a formal rank. As I have noted previously (Engel, 2001: 35), the rank of epifamily could be applied and the bees would be known as “Apoidae” (i.e., basing the name on the oldest, available family-group name: *vide infra*). This would only serve to confuse issues as this name is nearly identical to superfamily Apoidea (i.e., the currently employed suffix for the rank of epifamily is unfortunate and serious consideration should be given to changing it). I believe it most prudent to not employ the rank of epifamily and to instead use a non-ranked name for the bees. The utility of the name Aculeata, long used without a formal rank, is an excellent example of a significant lineage to which a non-ranked name has been applied successfully. Indeed, some recent authors have chosen to use the non-ranked name Apiformes for the bees. Apiformes was first introduced by Rayment (1935) for what is effectively the corbiculate apines of today and was later expanded to include all bees by Brothers (1975: 640). This name has, fortunately, not achieved universal usage and for myriad reasons I suggest that it not be adopted. Perhaps the longest used name for the bees has been Anthophila as proposed by

Latreille (1804: originally in the plural as *Anthophili*, but subsequently emended: *vide etiam* Latreille, 1805, 1807). This name was used for the bees as a whole by numerous authors throughout much of the 19th and early 20th centuries (e.g., Westwood, 1840; Smith, 1853, 1855; Cresson, 1887; Dalla Torre, 1896; Handlirsch, 1925). To the best of my knowledge, this is the oldest, formalized name applied to the bees as a whole and it is, moreover, quite descriptive (meaning “lovers of flowers”). I, therefore, recommend that this name be readopted for the bees, without formal rank (*vide* appendix 1).

FORMAT

The format for entries generally follows that employed in similar family-group name treatments (e.g., Engel and Krishna, 2004). Daggers (†) indicate names proposed for fossil lineages. When the author originally used an incorrect spelling for the stem (*nomen imperfectum*) I have provided the name as it first appeared, followed by how the name should have appeared with a corrected stem. For example, Cockerell and Atkins (1902) proposed the name *Ericrocinae* based on the type genus *Ericrocis*; however, the stem is incorrectly formed and should have been *Ericrocidinae*. Such incorrect original stems are denoted by “*recte* *Ericrocidinae*” and simply indicate how the name *should* have appeared. If the author simply employed a non standard suffix I have not considered these as incorrectly formed and therefore did not provide such corrections. For example, Lepeletier de Saint Fargeau (1841) proposed *Rhathymites*, based on *Rhathymus*, which has as its stem *Rhathym-* and the family-group name is therefore not corrected simply because he used a different suffix for the rank of family. Alternatively, in the same paper Lepeletier de Saint Fargeau (1841) proposed *Eulemites* based on *Eulaema*, which is corrected to *Eulaemites*, denoted as “*recte* *Eulaemites*”, since the stem is *Eulaem-*. The correct combining stem for any family-group name based on the name today is provided in each entry.

I have included in the list six *nomina nuda* that are based on available genus-group names. These names were not subsequently

made available in later publications (unlike some other family-group *nomina nuda*), appearing only as *nomina nuda*. Unlike some of the other names, there is no reason to validate these names, as they would be best considered synonyms of already available names (*vide* appendix 1).

As in my other papers, I have placed as the first date for each name and article the date as it is printed in publication. If the date for nomenclatorial purposes differs from the printed date (e.g., the publication appeared later than as printed, or the name is attributed to a different date by the ICZN for purposes of priority), then this date appears in brackets following the printed date. This format is also employed in the bibliography.

FAMILY-GROUP NAMES

1. *Andrenetae* Latreille, 1802a: 422. Type genus: *Andrena* Fabricius, 1775. Combining stem: *Andren-*. Note: *vide etiam* *Apiariae* (*vide infra*).

2. *Nomades* Latreille, 1802a: 425. Type genus: *Nomada* Scopoli, 1770. Combining stem: *Nomad-*. Note: *vide etiam* *Apiariae* (*vide infra*).

3. *Apiariae* Latreille, 1802a: 425. Type genus: *Apis* Linnaeus, 1758. Combining stem: *Ap-*. Notes: Although the family-group names based on *Andrena* and *Nomada* have page priority (or, in the case of *Nomades*, which appears on the same page, priority in position on the printed page itself) over that based on *Apis*, it is up to the first reviser to select which name is to be given nomenclatorial priority. Such priority has been universally given to the name based on *Apis* by entomologists since the earliest days of zoological nomenclature. As to competitions of priority between *Nomadidae* and *Andrenidae*, this seems to be of little concern, but I believe priority would be owed to *Andrenidae* should any such question arise owing to the usage of *Andrenidae* and *Apidae* as the main divisions of bees in many 19th century classifications.

There had been confusion in the past as to whether the name based on *Sphex* or on *Apis* had priority and should be used to form the superfamilial name encompassing both bees and digger wasps. Unlike the preceding

names (i.e., Andrenetae and Nomades), Sphegimae Latreille (1802b: 331) did not appear until a subsequent publication in the same year (November 1802, *vide* Dupuis, 1986) and is automatically junior to the three family-group names provided in Latreille (1802a), which appeared prior to April 1802. Apoidea takes priority over Sphecoidea (as would Andrenoidea and Nomadoidea) (*vide etiam* Michener, 1986a, and Menke, 1997).

4. Eucerae Latreille, 1802b: 376. Type genus: *Eucera* Scopoli, 1770. Combining stem: Eucer-.

5. Podalirii Latreille, 1802b: 377. Type genus: *Podalirius* Latreille, 1802a. Combining stem: Podaliri-.

6. Xylocopae Latreille, 1802b: 379. Type genus: *Xylocopa* Latreille, 1802b. Combining stem: Xylocop-.

7. Ceratinae Latreille, 1802b: 380. Type genus: *Ceratina* Latreille, 1802b. Combining stem: Ceratin-.

8. Megachiles Latreille, 1802b: 381. Type genus: *Megachile* Latreille, 1802a. Combining stem: Megachil-.

9. Euglossae Latreille, 1802b: 384. Type genus: *Euglossa* Latreille, 1802a. Combining stem: Eugloss-.

10. Bombi Latreille, 1802b: 385. Type genus: *Bombus* Latreille, 1802a. Combining stem: Bomb-.

11. Prosopiariae Fallén, 1813: 32. Type genus: *Prosopis* Fabricius, 1804 (*nec Prosopis* Jurine *In* Panzer, 1801, *nomen rejiciendum* [ICZN, 1939]). Combining stem: Prosopid-.

12. Panurgida Leach, 1815: 156. Type genus: *Panurgus* Panzer, 1806. Combining stem: Panurg-.

13. Anthrenoides Burmeister, 1829: 30, *nomen imperfectum* [*recte* Anthrenides]. Type genus: *Anthrena* Illiger, 1801, *nomen vanum*. Combining stem: Anthren-. Note: This name should not be confused with the genus-group name *Anthrenoides* DUCKE.

14. Melliturgites Newman, 1834: 379, *nomen imperfectum* [*recte* Melitturgites]. Type genus: *Melitturga* Latreille, 1809. Combining stem: Melitturg-.

15. Lithurgites Newman, 1834: 379. Type genus: *Lithurgus* Berthold, 1827. Combining stem: Lithurg-.

16. Apathites Newman, 1834: 379. Type

genus: *Apathus* Newman, 1834. Combining stem: Apath-.

17. Osmiites Newman, 1834: 401. Type genus: *Osmia* Panzer, 1806. Combining stem: Osmi-.

18. Anthophorini Dahlbom, 1835: 5. Type genus: *Anthophora* Latreille, 1803. Combining stem: Anthophor-.

19. Meliponites Lepeletier de Saint Fargeau, 1836: 407. Type genus: *Melipona* Illiger, 1806. Combining stem: Melipon-.

20. Chelostomidae Kirby, 1837: 270. Type genus: *Chelostoma* Latreille, 1809. Combining stem: Chelostom-.

21. Melectides Westwood, 1840 [1839]: 270. Type genus: *Melecta* Latreille, 1802a. Combining stem: Melect-.

22. Eulemites Lepeletier de Saint Fargeau, 1841: 218, *nomen imperfectum* [*recte* Eulaemites]. Type genus: *Eulaema* Lepeletier de Saint Fargeau, 1841. Combining stem: Eulaem-.

23. Colletides Lepeletier de Saint Fargeau, 1841: 295. Type genus: *Colletes* Latreille, 1802a. Combining stem: Collet-. Note: Although employed by Lepeletier de Saint Fargeau (1841) as a French vernacular, the name was adopted by later authors in latinized form and considered valid. Lepeletier de Saint Fargeau's name is, therefore, available as a family-group name, dating from the first publication employing it as a vernacular (ICZN, 1999a: Art. 11.7.2). The same condition holds true for other family-group names proposed by this author.

24. Psithyrides Lepeletier de Saint Fargeau, 1841: 423. Type genus: *Psithyrus* Lepeletier de Saint Fargeau, 1832. Combining stem: Psithyr-.

25. Philereimides Lepeletier de Saint Fargeau, 1841: 510. Type genus: *Phileremus* Latreille, 1809. Combining stem: Philereim-.

26. Rhathymites Lepeletier de Saint Fargeau, 1841: 539. Type genus: *Rhathymus* Lepeletier de Saint Fargeau and Audinet-Serville, 1828. Combining stem: Rhathym-.

27. Melittidae Schenck, 1860: 136. Type genus: *Melitta* Kirby, 1802. Combining stem: Melitt-.

28. Stelidae Schenck, 1860: 141, *nomen imperfectum* [*recte* Stelididae]. Type genus: *Stelis* Panzer, 1806. Combining stem: Stelid-.

29. Rhophitidae Schenck, 1866 [1867]:

322, *nomen imperfectum* [recte Rophitidae]. Type genus: *Rophites* Spinola, 1808. Combining stem: Rophit-.

30. Sphecodidae Schenck, 1868 [1869]: 316. Type genus: *Sphecodes* Latreille, 1804. Combining stem: Sphecod-.

31. Megillina Thomson, 1869: 7. Type genus: *Megilla* Fabricius, 1804. Combining stem: Megill-.

32. Halictina Thomson, 1869: 8. Type genus: *Halictus* Latreille, 1804. Combining stem: Halict-.

33. Coelioxynae Dalla Torre and Friese, 1894: 35. Type genus: *Coelioxys* Latreille, 1809. Combining stem: Coelioxys-.

34. Neolarrini Fox, 1894 [1895]: 302. Type genus: *Neolarra* Ashmead, 1890. Combining stem: Neolarr-.

35. Oxaeinae Ashmead, 1899a: 70. Type genus: *Oxaea* Klug, 1807a. Combining stem: Oxae-.

36. Anthidiinae Ashmead, 1899a: 72. Type genus: *Anthidium* Fabricius, 1804. Combining stem: Anthidi-.

37. Centrini Cockerell and Cockerell, 1901: 47, *nomen imperfectum* [recte Centridini]. Type genus: *Centris* Fabricius, 1804. Combining stem: Centrid-.

38. Melissodinae Robertson, 1901: 231. Type genus: *Melissodes* Latreille, 1829. Combining stem: Melissod-. Note: This name has been overlooked by most bee systematists. It was first noticed by M. G. Rightmyer and I am grateful that she has brought it to my attention.

39. Ericrocinae Cockerell and Atkins, 1902: 46, *nomen imperfectum* [recte Ericrocidae]. Type genus: *Ericrocis* Cresson, 1887. Combining stem: Ericrocid-.

40. Dioxinae Cockerell, 1902: 233, *nomen imperfectum* [recte Dioxyinae]. Type genus: *Dioxys* Lepeletier de Saint Fargeau and Audinet-Serville, 1825. Combining stem: Dioxy-.

41. Allodapinae Cockerell, 1902: 233. Type genus: *Allodape* Lepeletier de Saint Fargeau and Audinet-Serville, 1825. Combining stem: Allodap-.

42. Trypetini Robertson, 1903a: 164. *Nomen praeoccupatum* (nec Trypetidae Loew, 1861 [Diptera: Tephritidae]). Type genus: *Trypetes* Schenck, 1859 [1861], *nomen praeoccupatum* (nec *Trypetes* Schönherr,

1836 [Coleoptera: Curculionidae]). Combining stem: Trypet-. Note: The family-group name is also homonymous with a family-group name based on *Trypetes* Schönherr, 1836; i.e., Trypetinae Lacordaire, 1866 (originally spelled Trypetides by Lacordaire, but given a standard family-group suffix by Pascoe, 1870). The coleopterous taxon was emended to Trypetidinae (Pierce, 1919; ICZN, 1974), preserving Trypetinae for a lineage of flies based on the genus *Trypeta* Meigen, 1803. The family-group name based on *Trypetes* Robertson, 1903a cannot be conserved since its type genus is also a junior homonym (ICZN, 1999a: Art. 39).

43. Epeolinae Robertson, 1903b: 284. Type genus: *Epeolus* Latreille, 1802a. Combining stem: Epeol-.

44. Trachusinae Robertson, 1904: 37. Type genus: *Trachusa* Panzer, 1804b. Combining stem: Trachus-.

45. Exoneuridae Robertson, 1904: 37. Type genus: *Exoneura* Smith, 1854. Combining stem: Exoneur-.

46. Protandreninae Robertson, 1904: 38. Type genus: *Protandrena* Cockerell, 1896. Combining stem: Protandren-.

47. Dufoureae Robertson, 1904: 42. Type genus: *Dufourea* Lepeletier de Saint Fargeau, 1841. Combining stem: Dufoure-.

48. Nomiidae Robertson, 1904: 42. Type genus: *Nomia* Latreille, 1804. Combining stem: Nomi-.

49. Macropididae Robertson, 1904: 42. Type genus: *Macropis* Panzer, 1809. Combining stem: Macropid-.

50. Emphoridae Robertson, 1904: 43. Type genus: *Emphor* Patton, 1879. Combining stem: Emphor-.

51. Entechniini Cockerell, 1906: 99. Type genus: *Entechnia* Patton, 1879. Combining stem: Entechni-.

52. Palaeorhizidae Perkins, 1908: 29. Type genus: *Palaeorhiza* Perkins, 1908. Combining stem: Palaeorhiz-.

53. Meroglossidae Perkins, 1908: 29. Type genus: *Meroglossa* Smith, 1853. Combining stem: Merogloss-. Note: Although proposed tentatively by Perkins (1908) the name dates from that publication since prior to 1931 the simple formation of any family-group name based on an available genus-group name is

to be considered available (ICZN, 1999a: Art. 12.2.4).

54. Exaeretinae Cockerell, 1908: 41. Type genus: *Exaerete* Hoffmannsegg, 1817. Combining stem: Exaeret-.

55. Exomalopses Vachal, 1909: 7. Type genus: *Exomalopsis* Spinola, 1853. Combining stem: Exomalops-.

56. Melitomatae Vachal, 1909: 7. Type genus: *Melitoma* Lepeletier de Saint Fargeau and Audinet-Serville, 1828. Combining stem: Melitom-.

57. Diphaglossinae Vachal, 1909: 33. Type genus: *Diphaglossa* Spinola, 1851. Combining stem: Diphagloss-.

58. †Glyptapinae Cockerell, 1909b: 13. Type genus: *Glyptapis* Cockerell, 1909a. Combining stem: Glyptap-.

59. Oxystoglossini Schrottky, 1909a: 482. Type genus: *Oxystoglossa* Smith, 1853. Combining stem: Oxystogloss-.

60. Eucondylopsinae Friese, 1909: 99. Type genus: *Eucondylops* Brauns, 1902. Combining stem: Eucondylops-.

61. Hemisiinae Cockerell and Robbins, 1910: 183. Type genus: *Hemisia* Klug, 1807b. Combining stem: Hemisi-.

62. Gastrohalictinae Schrottky, 1911: 84. Type genus: *Gastrohalictus* Ducke, 1902. Combining stem: Gastrohalict-.

63. Epicharitina Schrottky, 1913: 259. Type genus: *Epicharis* Klug, 1807b. Combining stem: Epicharit-.

64. Tetraloniinae Schrottky, 1913: 263. Type genus: *Tetralonia* Spinola, 1838. Combining stem: Tetraloni-.

65. Hylaeidae Viereck, 1916: 737. Type genus: *Hylaeus* Fabricius, 1793. Combining stem: Hylae-. Note: The name Prosopiaridae Fallén, 1813 has priority over Hylaeidae; however, the family-group name based on *Hylaeus* has been conserved over that based on *Prosopis* (ICZN, 1993).

66. Dasypodinae Börner, 1919: 180 (Dasypodinae, *nomen emendatum* [ICZN, 1999b]). Type genus: *Dasypoda* Latreille, 1802a. Combining stem: Dasypoda-. Note: The stem of the family-group name based on *Dasypoda* Latreille, 1802a was altered from Dasypod- to Dasypoda- by ICZN (1999b).

67. Halictoidinae Börner, 1919: 181. Type genus: *Halictoides* Nylander, 1848. Combining stem: Halictoid-.

68. Nomioidini Börner, 1919: 181. Type genus: *Nomioides* Schenck, 1866 [1867]. Combining stem: Nomioid-.

69. Bremidae Frison, 1919: 157. Type genus: *Bremus* Panzer, 1804a (*nec Prosopis* Jurine in Panzer, 1801, *nomen rejiciendum* [ICZN 1939: Op. 135]). Combining stem: Brem-.

70. Perditinae Robertson, 1922: 160. Type genus: *Perdita* Smith, 1853. Combining stem: Perdit-.

71. Calliopsinae Robertson, 1922: 160. Type genus: *Calliopsis* Smith, 1853. Combining stem: Calliops-.

72. Anthemurginae Robertson, 1922: 160. Type genus: *Anthemurgus* Robertson, 1902a. Combining stem: Anthemurg-.

73. Systrophini Handlirsch, 1925: 819. Type genus: *Systropha* Illiger, 1806. Combining stem: Systroph-.

74. Odyneropsini Handlirsch, 1925: 821. Type genus: *Odyneropsis* Schrottky, 1902. Combining stem: Odynerops-.

75. Osirini Handlirsch, 1925: 821. Type genus: *Osiris* Smith, 1854. Combining stem: Osir-.

76. Ammobatini Handlirsch, 1925: 821. Type genus: *Ammobates* Latreille, 1809. Combining stem: Ammobat-.

77. Melanempidini Handlirsch, 1925: 821. Type genus: *Melanempis* Saussure, 1890 [1892]. Combining stem: Melanempid-.

78. Chrysanthedinae Handlirsch, 1925: 821. Type genus: *Chrysantheda* Perty, 1833. Combining stem: Chrysanthed-.

79. Aglainae Handlirsch, 1925: 822. Type genus: *Aglae* Lepeletier de Saint Fargeau and Audinet-Serville, 1825. Combining stem: Agla-.

80. Augochloridae Beebe, 1925: 102. Type genus: *Augochlora* Smith, 1853. Combining stem: Augochlor-. Note: This name is widely indicated to date from Moure (1943). However, as noted by Engel (1999a, 2000), the correct author and date is Beebe (1925).

81. Lasioglossinae Robertson, 1926: 117. Type genus: *Lasioglossum* Curtis, 1833. Combining stem: Lasiogloss-.

82. Chloralictini Robertson, 1926: 117. Type genus: *Chloralictus* Robertson, 1902b. Combining stem: Chloralict-.

83. Pasititidae Robertson, 1926: 118, *nomen imperfectum* [recte Pasitidae]. Type ge-

nus: *Pasites* Jurine, 1807. Combining stem: Pasit-.

84. Xeromelissinae Cockerell, 1926: 222. Type genus: *Xeromelissa* Cockerell, 1926. Combining stem: Xeromeliss-.

85. Hylaeoidinae Cockerell, 1930a: 148, *nomen imperfectum* [recte Hyleoidinae]. Type genus: *Hyleoides* Smith, 1853. Combining stem: Hyleoid-.

86. Neopasiphaeinae Cockerell, 1930a: 148. Type genus: *Neopasiphae* Perkins, 1912. Combining stem: Neopasiphae-.

87. Ctenoplectridae Cockerell, 1930a: 148. Type genus: *Ctenoplectra* Kirby in Kirby and Spence, 1826. Combining stem: Ctenoplectr-.

88. Fideliidae Cockerell, 1932: 358. Type genus: *Fidelia* Friese, 1899. Combining stem: Fideli-.

89. Stenotritini Cockerell, 1934: 6. Type genus: *Stenotritus* Smith, 1853. Combining stem: Stenotrit-. Note: The family-group names established by Cockerell in this account of Australian bees (namely, Stenotritini, Phenacolletini, and Paracolletini) are quite problematic, although the solution to the nomenclatorial dilemma proves to be merely academic as it does not alter their usage. None of the three family-group names proposed in this article are provided with a description and therefore fail one of the criteria for availability of such names established after 1930 (ICZN, 1999a: Art. 13.1). Cockerell merely notes at the top of page 6 in his article, "The genera may be divided into three tribes, Stenotritini, Phenacolletini and Paracolletini", and then later includes them as headers for sections considering the species in further detail (but again, does not accompany the tribal headings with any characters purported to differentiate them from each other or from other such suprageneric groups). Thus, Stenotritini, Phenacolletini, and Paracolletini should be considered *nomina nuda*. However, while Cockerell's names fail Article 13.1, two of them (i.e., Stenotritini and Paracolletini) satisfy the provision to 13.1 provided by 13.2.1 (ICZN, 1999a) whereby a name proposed after 1930 and before 1961 that lacks a description is to be considered available from its original publication *only* if it was employed as valid before 2000 and was not rejected by any author applying Article 13.1 between 1960 and

2000. Thus, the usage of Stenotritini and Paracolletini by subsequent authors (e.g., Michener, 1944, 1965) and the fact that no author rejected the names as *nomina nuda* between 1960 and 2000 permits these two names to be considered available from Cockerell (1934). This is not the case for Phenacolletini (*vide infra*).

90. Phenacolletini Cockerell, 1934: 7, *nomen nudum*. Type genus: *Phenacolletes* Cockerell, 1905. Combining stem: Phenacollet-. Note: As discussed under Stenotritini (*vide supra*), Phenacolletini was technically a *nomen nudum* in Cockerell (1934). Unlike Stenotritini and Paracolletini, which are conserved by the provisions of Article 13.2.1 (ICZN, 1999a), Phenacolletini was not employed as valid by later authors and thus remains unavailable. While it might be argued that Rayment (1935, p. 186) subsequently employed the name Phenacolletini, I cannot consider this to be true since Rayment is merely quoting the line from Cockerell (1934). I cannot see any reason why Phenacolletini should be made available herein as it is clearly a synonym.

91. Paracolletini Cockerell, 1934: 8. Type genus: *Paracolletes* Smith, 1853. Combining stem: Paracollet-. Note: Refer to discussion under Stenotritini (*vide supra*).

92. Biastini Linsley and Michener, 1939: 272. Type genus: *Biastes* Panzer, 1806. Combining stem: Biast-.

93. Neopasitini Linsley and Michener, 1939: 274. Type genus: *Neopasites* Ashmead, 1898. Combining stem: Neopasit-.

94. Epeoloidini Linsley and Michener, 1939: 294. Type genus: *Epeoloides* Giraud, 1863. Combining stem: Epeoloid-.

95. Protepeolini Linsley and Michener, 1939: 301. Type genus: *Protepeolus* Linsley and Michener, 1937. Combining stem: Protepeol-.

96. Heriadini Michener, 1941: 152. Type genus: *Heriades* Spinola, 1808. Combining stem: Heriad-.

97. Augochloropsini Moure, 1943: 462. Type genus: *Augochloropsis* Cockerell, 1897. Combining stem: Augochlorops-. Note: Although a character description is not provided for this tribe, Moure's statement that it is proposed for Vachal's (1911) "*Halicti vibrissati*" can be considered biblio-

graphic reference to a description, therefore making the name available in this publication by indication.

98. Megaloptinae Moure, 1943: 479. Type genus: *Megalopta* Smith, 1853. Combining stem: Megalopt-.

99. Caupolicanini Michener, 1944: 238. Type genus: *Caupolicana* Spinola, 1851. Combining stem: Caupolican-.

100. Euryglossinae Michener, 1944: 238. Type genus: *Euryglossa* Smith, 1853. Combining stem: Eurygloss-.

101. Chilicolinae Michener, 1944: 240. Type genus: *Chilicola* Spinola, 1851. Combining stem: Chilicol-.

102. Ancylini Michener, 1944: 273. Type genus: *Ancyla* Lepeletier de Saint Fargeau, 1841. Combining stem: Ancy-.

103. Caenoprosopidini Michener, 1944: 275. Type genus: *Caenoprosopis* Holmberg, 1886. Combining stem: Caenoprosopid-.

104. Townsendiellini Michener, 1944: 277. Type genus: *Townsendiella* Crawford, 1916. Combining stem: Townsendiell-.

105. Ammobatoidini Michener, 1944: 277. Type genus: *Ammobatoides* Radoszkowski, 1868 [1867]. Combining stem: Ammobatoid-.

106. Eufrieseini Moure, 1944: 12. Type genus: *Eufriesea* Cockerell, 1908a. Combining stem: Eufriese-.

107. Eulonchopriinae Moure, 1945: 140. Type genus: *Eulonchopria* Brèthes, 1909. Combining stem: Eulonchopri-.

108. Lonchopriinae Moure, 1945: 141. Type genus: *Lonchopria* Vachal, 1905b. Combining stem: Lonchopri-.

109. Dissoglottini Moure, 1945: 142. Type genus: *Dissoglotta* Moure, 1945. Combining stem: Dissoglott-.

110. Trigonini Moure, 1946: 611. Type genus: *Trigona* Jurine, 1807. Combining stem: Trigon-.

111. Lestrimelittini Moure, 1946: 611. Type genus: *Lestrimelitta* Friese, 1903b. Combining stem: Lestrimelitt-.

112. Dianthidiini Moure, 1947: 15. Type genus: *Dianthidium* Cockerell, 1900. Combining stem: Dianthidi-.

113. Trichothurgini Moure, 1949: 268. Type genus: *Trichothurgus* Moure, 1949. Combining stem: Trichothurg-.

114. Pararhophitini Popov, 1949: 507.

Type genus: *Pararhophites* Friese, 1898. Combining stem: Pararhophit-.

115. Euherbstiinae Moure, 1950: 303. Type genus: *Euherbstia* Friese, 1925. Combining stem: Euherbsti-.

116. Ptiloglossidiini Moure, 1953: 71. Type genus: *Ptiloglossidia* Moure, 1953. Combining stem: Ptiloglossidi-.

117. Caenohalictinae Michener, 1954: 97. Type genus: *Caenohalictus* Cameron, 1903. Combining stem: Caenohalict-.

118. Canephorulini Michener, LaBerge, and Moure, 1955: 207. Type genus: *Canephorula* Jörgensen, 1909. Combining stem: Canephorul-.

119. Eucerinodini Michener and Moure, 1957: 444. Type genus: *Eucerinoda* Michener and Moure, 1957. Combining stem: Eucerinod-.

120. Tetrapediini Michener and Moure, 1957: 442. Type genus: *Tetrapedia* Klug, 1810. Combining stem: Tetrapedi-.

121. Ctenioschelini Michener, 1965a: 10, *nomen nudum*. Type genus: *Ctenioschelus* Romand, 1840. Combining stem: Ctenioschel-. Note: This name is not available from this publication since it was merely included in a list of tribes but not accompanied by a description as required for family-group names after 1930 (ICZN, 1999a: Arts. 13.1, 13.2). It has not been subsequently made available and I do not believe it is necessary to do so now as it is assuredly a synonym of *Ericrocidini*.

122. Mydrosomini Michener, 1966: 719, *nomen imperfectum* [*recte* Mydrosomatini]. Type genus: *Mydrosoma* Smith, 1879. Combining stem: Mydrosomat-.

123. Holcopasitini Rozen, 1966: 30. Type genus: *Holcopasites* Ashmead, 1899a. Combining stem: Holcopasit-.

124. Pachyanthidiini Pasteels, 1969: 6. Type genus: *Pachyanthidium* Friese, 1905. Combining stem: Pachyanthidi-.

125. Icteranthidiini Pasteels, 1969: 7. Type genus: *Icteranthidium* Michener, 1948. Combining stem: Icteranthidi-.

126. Euaspini Pasteels, 1969: 7. Type genus: *Euaspis* Gerstcker, 1857. Combining stem: Euasp-.

127. Thrinchostomini Sakagami, 1974: 258, *nomen imperfectum* [*recte* Thrinchostomatini]. Type genus: *Thrinchostoma* Saus-

sure, 1890 [1892]. Combining stem: Thrinchostomat-.

128. Habropodini Marikovskaya, 1976: 688. Type genus: *Habropoda* Smith, 1854. Combining stem: Habropod-.

129. Isepeolini Rozen, Eickwort, and Eickwort, 1978: 18. Type genus: *Isepeolus* Cockerell, 1907. Combining stem: Isepeol-.

130. Meganomiinae Michener, 1981: 18. Type genus: *Meganomia* Cockerell, 1909c. Combining stem: Meganomi-.

131. Promelittini Michener, 1981: 60. Type genus: *Promelitta* Warncke, 1977. Combining stem: Promelitt-.

132. Sambini Michener, 1981: 60. Type genus: *Samba* Friese, 1908. Combining stem: Samb-.

133. Manueliini Sakagami and Michener, 1987: 439. Type genus: *Manuelia* Vachal, 1905a. Combining stem: Manueli-.

134. Protomeliturgini Ruz, 1991: 215. Type genus: *Protomeliturga* Ducke, 1912. Combining stem: Protomeliturg-.

135. Brachynomadini Roig-Alsina and Michener, 1993: 157. Type genus: *Brachynomada* Holmberg, 1886. Combining stem: Brachynomad-. Note: Descriptions, which are required after 1930 (ICZN, 1999a: Arts. 13.1., 13.2, “must be accompanied by a description that states in words characters that are purported to differentiate the taxon”; and were also required by the edition of the Code that was applicable in 1993; namely, ICZN, 1985: Art. 13a), of new family-group taxa are difficult to discern in this publication and I do not believe that the extraction of characters mapped from their numerous cladograms can be considered diagnoses since no unique tree or suite of characters are explicitly cited for defining their new taxa. However, family-group names can be made available by reference, and in the case of Brachynomadini these authors explicitly refer to the definition of the “melanomadine complex of Alexander (1990)”. I, therefore, believe them to have satisfactorily met the criteria for availability by bibliographic reference to a description.

136. Tapinotaspidini Moure, 1992 [1994]: 306. Type genus: *Tapinotaspis* Holmberg, 1903. Combining stem: Tapinotaspid-. Note: This name was first mentioned in Roig-Alsina and Michener (1993: 159), as Tapino-

taspini Roig-Alsina and Michener, 1993, *nomen imperfectum* [recte Tapinotaspidini] and is widely held to be available from that publication (e.g., Michener, 1997). For the same reasons described for Hexepeolini (*vide infra*) I believe that Tapinotaspidini was a *nomen nudum* in 1993 as no description was provided as clearly required by the Code (ICZN, 1999a: Arts. 13.1, 13.2; and was also required by the edition of the Code which was applicable in 1993; namely, ICZN, 1985: Art. 13a). It might be believed that the statement, “These genera are those of sections 1, 2, and 5 of Exomalopsini as understood by Michener and Moure (1957)”, could qualify as a bibliographic reference, but there are no diagnostic traits given to unite groups 1, 2, and 5 together in the cited paper (which would, therefore, represent characters of Tapinotaspidini). As such, the reference fails to validate the name by indication. However, Moure’s (1992: NB: this paper dates for nomenclatorial purposes from 1994) subsequent account of the tribe includes a description and employs a family-group name based on an available genus-group name. I, therefore, consider the name as having been first made available in the latter paper, despite Michener’s (1997) comments to the contrary (Roig-Alsina, 1997, also includes what can be considered a description of the tribe). Since both Tapinotaspidini and Paratetrapediini date from the same paper and are synonyms, I here select the former name for the tribe following current taxonomic usage.

137. Paratetrapediini Moure, 1992 [1994]: 306. Type genus: *Paratetrapedia* Moure, 1941. Combining stem: Paratetrapedi-.

138. Hexepeolini Rozen, 1996: 188. Type genus: *Hexepeolus* Linsley and Michener, 1937. Combining stem: Hexepeol-. Note: This name suffers the same problem as described for Brachynomadini and Tapinotaspidini (*vide supra*). The name was first mentioned in Roig-Alsina and Michener (1993: 157) as Hexepeolini Roig-Alsina and Michener, 1993, and is widely held to be available from that publication. However, I believe that Hexepeolini was a *nomen nudum* in 1993. I cannot conscientiously consider it as available from Roig-Alsina and Michener (1993) despite the inclusion of only *Hexepeolus* since

no description or bibliographic reference to such a description is provided (as explicitly required by the Code [ICZN, 1999a: Arts. 13.1, 13.2] and was also required by the edition of the Code that was applicable in 1993; namely, ICZN, 1985: Art. 13a). Inclusion of a single genus does not validate the name, as such an indication is not valid after 1930. Thus, I consider Hexepeolini Roig-Alsina and Michener, 1993 a *nomen nudum* and unavailable from that work. The next paper providing a clear description that can loosely be attributed to the tribe (i.e., providing in words characters that purport to differentiate the taxon from others), albeit based on larval characters, is that of Rozen (1996) who, prior to his description, states, “was assigned to its own monotypic tribe, the Hexepeolini” (NB: prior to 1999 it was not necessary to explicitly state that a family-group name was new). Since Rozen (1996) uses a family-group name based on an available genus-group name and provides a differential diagnosis that can be loosely attributed to the tribe, I consider the name Hexepeolini to have been made available in that work.

139. Teratognathini Silveira, 1995: 449. Type genus: *Teratognatha* Ogloblin, 1956. Teratognath-.

140. †Electrapina Engel, 1998a: 99. Type genus: †*Electrapis* Cockerell, 1908b. Combining stem: Electrap-.

141. Camptopoeumini Patiny, 1999: 270, *nomen imperfectum* [recte Camptopoeini]. Type genus: *Camptopoeum* Spinola, 1843. Combining stem: Camptopoe-. Note: *Camptopoeum* is a combination of the Greek words *kampto* (meaning “bend”) and *poieo* (meaning “make”). The latter word, when latinized, has as its stem poe- and, as such, the spelling of the family-group name is corrected herein.

142. Mermiglossini Patiny, 1999: 270. Type genus: *Mermiglossa* Friese, 1912. Combining stem: Mermigloss-.

143. Panurginini Patiny, 1999: 271. Type genus: *Panurginus* Nylander, 1848. Combining stem: Panurgin-.

144. †Chalicodomopsini Engel, 1999b: 4, *nomen nudum*. Type genus: †*Chalicodomopsis* Engel, 1999b. Combining stem: Chalicodomops-. Note: This family-group name is unavailable, since after 1930 all fam-

ily-group names require a formal description (ICZN, 1999a: Art. 13.2). There is no reason to validate it herein.

145. Corynurina Engel, 2000: 51. Type genus: *Corynura* Spinola, 1851. Combining stem: Corynur-.

146. Ischnomelissiti Engel, 2000: 66. Type genus: *Ischnomelissa* Engel, 1997. Combining stem: Ischnomeliss-.

147. Thectochloriti Engel, 2000: 66. Type genus: *Thectochlora* Moure, 1940. Combining stem: Thectochlor-.

148. Megaloptidiiti Engel, 2000: 66. Type genus: *Megaloptidia* Cockerell, 1900. Combining stem: Megaloptidi-.

149. Agapostemonina Engel, 2000: 67. Type genus: *Agapostemon* Guérin-Ménéville, 1844. Combining stem: Agapostemon-.

150. Alocandreninae Michener, 2000: 228. Type genus: *Alocandrena* Michener, 1986b. Combining stem: Alocandren-.

151. Liphanthini Michener, 2000: 262, *nomen nudum*. Type genus: *Liphanthus* Reed, 1894. Combining stem: Liphanth-. Note: This name was first used in an unpublished doctoral dissertation by Ruz (1987) and was thus unavailable. Its subsequent usage has not made it available and there seems little reason to do so since it would merely be a synonym of Protandrenini.

152. Austropanurgini Michener, 2000: 262, *nomen nudum*. Type genus: *Austropanurgus* Toro, 1980. Combining stem: Austropanurg-. Note: This name suffers the same problems as Liphanthini; see note for Liphanthini (*vide supra*).

153. Hoplitini Nobile and Tomarchio, 2000: 46, *nomen nudum*. Type genus: *Hoplitis* Klug, 1807b. Combining stem: Hoplit-.

154. Penapini Engel, 2001: 37. Type genus: *Penapis* Michener, 1965b. Combining stem: Penap-.

155. †Paleomelittidae Engel, 2001: 41. Type genus: †*Paleomelitta* Engel, 2001. Combining stem: Paleomelitt-.

156. Redivivini Engel, 2001: 45. Type genus: *Rediviva* Friese, 1911. Combining stem: Rediviv-.

157. †Eomacropidini Engel, 2001: 46. Type genus: †*Eomacropis* Engel, 2001. Combining stem: Eomacropid-.

158. †Protolithurgini Engel, 2001: 51.

Type genus: †*Protolithurgus* Engel, 2001. Combining stem: Protolithurg-.

159. †*Ctenoplectrellina* Engel, 2001: 54. Type genus: †*Ctenoplectrella* Cockerell, 1909a. Combining stem: Ctenoplectrell-.

160. †*Boreallodapini* Engel, 2001: 77. Type genus: †*Boreallodape* Engel, 2001. Combining stem: Boreallodap-.

161. †*Electrobombini* Engel, 2001: 88. Type genus: †*Electrobombus* Engel, 2001. Combining stem: Electrobomb-.

162. †*Melikertini* Engel, 2001: 112. Type genus: †*Melikertes* Engel, 1998a. Combining stem: Melikert-.

163. *Meliturgulini* Engel, 2001: 179. Type genus: *Meliturgula* Friese, 1903a. Combining stem: Meliturgul-. Note: *Meliturgulini* replaces the name *Paramelitturgini* Patiny, 1999: 272, *nomen illegitimum*, the latter of which was not based on an available genus-group name (and is, therefore, excluded from the list presented here).

164. *Nolanomelissini* Rozen and Ascher *in* Rozen, 2003: 102. Type genus: *Nolanomelissa* Rozen, 2003. Combining stem: Nolanomeliss-.

165. *Rhogepeolina* Rightmyer, 2004: 15. Type genus: *Rhogepeolus* Moure, 1955. Combining stem: Rhogepeol-.

166. *Thalestriina* Rightmyer, 2004: 17. Type genus: *Thalestria* Smith, 1854. Combining stem: Thalestri-.

167. *Neofideliina* Engel, 2004: 826. Type genus: *Neofidelia* Moure and Michener, 1955. Combining stem: Neofideli-.

168. *Scraptrinae* Ascher and Engel, herein (*vide infra*). Type genus: *Scrapter* Lepeletier de Saint Fargeau and Audinet-Serville, 1828. Combining stem: Scraptr-.

169. *Neffapini* Ascher, herein (*vide infra*). Type genus: *Neffapis* Ruz *in* Rozen and Ruz, 1995. Combining stem: Neffap-.

170. *Afrodasyopodini* Engel, herein (*vide infra*). Type genus: *Afrodasyopoda* Engel, herein (*vide infra*). Combining stem: Afrodasyopod-.

171. *Hesperapina* Ascher and Engel, herein (*vide infra*). Type genus: *Hesperapis* Cockerell, 1898. Combining stem: Hesperap-.

172. *Macrogaleina* Engel, herein (*vide infra*). Type genus: *Macrogalea* Cockerell, 1930b. Combining stem: Macrogale-.

173. *Ancyloscelidina* Engel and Michener,

herein (*vide infra*). Type genus: *Ancyloscelis* Latreille, 1829. Combining stem: Ancyloscelid-. Note: The name was first mentioned in Roig-Alsina and Michener (1993: 159), as *Ancyloscelina* Roig-Alsina and Michener, 1993, *nomen imperfectum* [*recte* Ancyloscelidina]. However, this name suffers the same problem as described for *Brachynomadini* and I believe that Ancyloscelidina was a *nomen nudum* in 1993. I cannot consciously consider it as available from Roig-Alsina and Michener (1993) despite the inclusion of only *Ancyloscelis* since no description or bibliographic reference to such a description is provided (as explicitly required by the Code [ICZN, 1999a: Arts. 13.1, 13.2] and was also required by the edition of the Code that was applicable in 1993; namely, ICZN, 1985: Art. 13a). Indeed, the subtribe is “diagnosed” by its exclusion from the subtribe Emphorina, which is not explicitly diagnosed therein either and thus one cannot extrapolate the characters purported to differentiate Ancyloscelidina. Inclusion of a single genus does not validate the name, as such an indication is not valid after 1930. Thus, I consider Ancyloscelidina Roig-Alsina and Michener, 1993 a *nomen nudum* and unavailable from that work. Unfortunately, no paper between 1993 and the publication of the most recent edition of the Code (ICZN, 1999a) has made the name available. While Michener (2000) also employed the subtribal name, post 1999 a family-group name must explicitly state that it is proposed as new, cite a type genus, and be accompanied by a description purporting characters that diagnose the taxon (ICZN, 1999a: Art. 16), the first two of which are not included therein. No such account for Ancyloscelidina has appeared and the name is, therefore, newly made available herein (*vide infra*).

SYSTEMATICS

Below are provided descriptions for six new family-group taxa. The placement of the new taxa among other bee lineages can be determined from the outline provided in appendix 1 as well as under the comments provided for each. In the diagnoses, the concepts of the various higher taxa of Apoidea are those as outlined in appendix 1 (e.g., Hyla-

einae herein is used in a broader sense to encompass the former subfamilies Euryglossinae and Xeromelissinae as tribes).

SCRAPTRINAE Ascher and Engel,
new subfamily

TYPE GENUS: *Scrapter* Lepeletier de Saint Fargeau and Audinet-Serville, 1828 (*nec* *Scrapter* Lepeletier de Saint Fargeau, 1841).

DIAGNOSIS: Body form of adults elongate as in many Xeromelissini; nonmetallic; vestiture sparse in most species. Premental fovea present (as in Xeromelissini and Hylaeini: homologous foveae are less well defined in Euryglossini; absent in true Paracolletinae, Colletinae *s. str.*, and Diphaglossinae); galeal comb reduced, usually to 3–4 small bristles. Facial foveae a narrow groove in many species, relatively broad in certain species but never densely setose and confined to a broad band between the compound eye and the lateral ocellus as in Colletinae *s. str.* and Diphaglossinae. Metabasitibial plate present (in many species carinate margin is interrupted, and in some species distinct lobes or tubercles are found resembling the tubercles characteristic of Euryglossini; absence of metabasitibial plates in Xeromelissini and Hylaeini may be correlated with cavity-nesting habits characteristic of these tribes; ground nesting in certain members of these tribes is likely secondary). Inner metatibial spur of female straight, tapering, ciliate; scopa of hind legs well developed; hind legs of males of certain species modified, with metafemur swollen and metabasitarsus expanded posteriorly towards apex (this modification is closely paralleled in many xeromelissine males: compare illustrations of *Scrapter* in Eardley, 1996, with those of xeromelissines in Toro and Moldenke, 1979). Forewing with two submarginal cells; second submarginal cell more than two-thirds as long as first (much shorter in most Hylaeinae *s. lato*, excepting, e.g., *Hyleoides*); pterostigma usually large, receiving r-rs towards middle. Pygidial and prepygidial fimbria of female present. Male S7 with apical lobes reduced or absent; if present, lobes small, aetose, laterally directed (Eardley, 1996: 37, twice erroneously referred to S8 of *Scapter* as “seventh sternite”); his diagnosis on the following page is

correct), elongate and with apex protruding externally in dorsal view, superficially resembling a pygidial plate, which is absent in *Scrapter* males (S8 internal in most other colletids). Mature larva with genal area laterally expanded; hypostomal ridge forming right angle with posterior thickening of head capsule; epistomal ridge arching dorsally to level of antennae; anterior tentorial pit extremely low in position (low anterior tentorial pits may be partially correlated with a strongly arched epistomal ridge); inner apical surface of maxilla rounded; salivary lips absent; integument of body not spiculate (McGinley, 1981); furthermore, all *Scrapter* species possess characters general to colletids that do not spin cocoons and lack many other distinctive larval and adult apomorphies unique to Diphaglossinae, Colletinae *s. str.*, Hylaeini, and Xeromelissini (e.g., McGinley, 1981; Michener, 2000; Ascher, 2004).

DISTRIBUTION: The 31 species of Scaptrinae are endemic to southern Africa (South Africa, Namibia, and Zimbabwe). Most are found in the semiarid Nama-Karoo and Succulent Karoo biomes (Eardley, 1996). True Paracolletinae are absent from Africa. Most species of the latter subfamily occur in temperate South America and Australia but a few derived taxa extend to adjacent areas such as North America north to southeastern Arizona and various islands near Australia such as Misoöl, New Guinea, Lord Howe, New Zealand, and New Caledonia (Michener, 2000).

COMMENTS: Presently the type genus is the only genus recognized in the subfamily. However, *Scrapter* is notably diverse and groups of its species differ by characters often used to delimit bee genera and subgenera (e.g., Michener, 2000). Once a cladistic study of the genus has been undertaken segregation of the species into subgenera would certainly appear warranted.

Scrapter has traditionally been placed in the Colletinae: Paracolletini (e.g., Michener, 1944), but neither the subfamilial nor the tribal placement has ever been supported by any substantial apomorphic characters, and Paracolletini has long been recognized as an exceptionally plesiomorphic and probably paraphyletic bee group (e.g., Michener, 1944,

1989). McGinley (1981: cladogram F) mapped his larval characters onto Michener's (1944) phylogenetic hypothesis. These supported the clade of Hylaeinae *s. lato* but otherwise provided little support for the traditional higher classification of the Colletidae. McGinley's (1981) preferred cladogram (cladogram D) for the paraphyletic Paracolletini demonstrated great divergence between *Scapter* and all other paracolletines, including the superficially similar *Callomelitta*. Cladograms including additional colletid lineages (e.g., his cladogram J) showed that *Scapter* is misplaced in Paracolletinae and demonstrated cladistic relationships among *Scapter*, *Colletes*, and Hylaeinae *s. lato*, excluding all paracolletines. Unfortunately, he did not propose a new classification based on his larval data.

Michener (2000) suggested that *Scapter* and *Callomelitta* are distinctive enough to warrant tribal status if separated from the old Paracolletini, but he presented far more compelling evidence demonstrating distinctiveness of the former as compared to the latter, and did not justify retaining *Scapter* at any rank within any existing subfamily. Michener (op. cit.) may be correct to interpret the tuberculate margin of the metabasitibial plate in certain *Scapter* and certain euryglossines as independently derived, but this parallelism is unusual and therefore suggestive of rather close relationship. His interpretation of similarities between the facial foveae of *Scapter* and those of *Callomelitta* and *Eulonchopria* proper as convergent is also likely correct, but similarity to the fovea of Hylaeini and Euryglossini may be truly homologous (e.g., Ascher, 2004). Silveira et al. (2002) recognized that *Scapter* did not belong in Paracolletinae, which they recognized as distinct from Colletinae *s. str.*, but did not propose a new subfamily for this taxon, which was extralimital to their study. Ascher (2004) presented support for a sister relationship between *Scapter* and the clade (Xeromelissini + (Euryglossini + Hylaeini)) (= Hylaeinae *s. lato* in the current classification: *vide* appendix 1) exclusive of Stenotritinae, Diphaglossinae, Paracolletinae, and Colletinae *s. str.*, in an analysis including adult and larval characters (notably those of McGinley, 1981).

Scaptrinae is one of six recognized subfamilies of Colletidae along with Stenotritinae, Diphaglossinae, Colletinae *s. str.* (including only *Colletes* and *Moureocotelles*), Paracolletinae (*sensu* Silveira et al., 2002, with *Scapter* formally excluded), and Hylaeinae (*sensu novum*, with the traditional subfamilies Euryglossinae and Xeromelissinae classified as tribes therein). Monophyly of each of the subfamilies Stenotritinae, Diphaglossinae, Colletinae *s. str.*, and Hylaeinae *s. lato* is supported by strong apomorphies (e.g., Michener, 2000), as is the monophyly of the tribes Hylaeini and Xeromelissini. Stenotritines lack colletid glossal synapomorphies and associated cellophane-like cell linings, which suggests placement basal to the remaining colletids, a position supported by some molecular data (e.g., Brady and Danforth, 2004). If shown to be sister to Colletidae, recognition of this taxon as a family (Michener, 2000) or subfamily (Engel, 2001, and herein) would both be consistent with phylogeny. Apomorphic character support for Euryglossini is more limited, but paraphyly with respect to Hylaeini has not been proposed. The status of Paracolletinae excluding *Scapter* as a monophyletic group is equivocal; there is no compelling evidence demonstrating either monophyly or paraphyly with respect to other subfamilies. In the absence of the latter evidence it is appropriate to retain the group pending further study. *Callomelitta* possesses many unusual characters (e.g., McGinley, 1981; Michener, 2000), but no strong evidence suggests that it is related to a nonparacolletine lineage.

NEFFAPINI Ascher, new tribe

TYPE GENUS: *Neffapis* Ruz in Rozen and Ruz, 1995.

DIAGNOSIS: (based, in part, upon description and illustrations in Rozen and Ruz, 1995). Glossa greatly elongate, each approximately four times as long as prementum, reaching beyond middle of metasoma in repose; glossal apex attenuate; labial palpus greatly elongate, three-segmented, first segment short, less than half as long as second segment, third segment extremely elongate, nearly four times as long as second; maxillary palpus minute, two-segmented; parag-

lossa short, widened distally; mandible with basal projection. Labrum flat; clypeus of female maculated with yellow (as in many other Panurgini, unlike most protandrenines). Antennal sockets unusually low on face (as in the otherwise dissimilar Calliopsini). Facial foveae shallow, poorly delimited (well-defined in most protandrenines, but also weak in many Panurgini *sensu* Ascher, 2004). Forewing with two submarginal cells (three in some protandrenines, Melitturgina, and Meliturgulina). Fovea of second metasomal tergum almost invisible (usually evident in protandrenines). S6 of female with inner surface laterad straight basal margin on each side with short, longitudinal sclerotized ridge (as in *Panurginus* and some other Old World panurgines, ridge extending across sternum in most protandrenines). S6 of male quadrate, apex very slightly emarginate; S7 of male with bilobed apex exposed; with short, subtriangular apical lobes broadly joined to broad body of sternum; basal apodemal arms broadly joined to sternal body (as in many Panurgini such as *Camptopoeumina*, unlike most protandrenines, which have smaller sternal bodies, elongate slender basal apodemal arms, and well-defined, membranous apical lobes); S8 of male narrowed basally to form spiculum (as in many Panurgini). Penis valves and penis fused in basal half; gonostylus angled orthogonal to gonocoxites; gonostylus slender and well-differentiated from much broader gonocoxites by membrane; volsellae free mesally. Mature larva with median section of epistomal ridge present but weak; antennal prominences low; maxillary apex (except for palpus) approximately in line with labial apex as seen in lateral view, labium therefore not greatly recessed relative to maxilla as in true protandrenines (Rozen and Ruz, 1995); thoracic tubercles unmodified as in most Panurgini, not forming lateral pockets between the prothorax and mesothorax and between the mesothorax and metathorax as in true Protandrenini and *Melitturga* (Rozen and Ruz, 1995; Rozen and Yanega, 1999). Pupa with few tubercles; mesoscutum, tegula, and base of metatibia on outer surface lacking distinct tubercles; terminal spine short, apically rounded, and not sclerotized (Rozen and Ruz, 1995).

DISTRIBUTION: The single species of *Neffapis* is endemic to the Coquimban desert of central Chile (Region IV, the Coquimban Region) at the southern end of the Atacama Desert, where it occurs with another Coquimban endemic andrenid genus, *Euherbstia* (vide Rozen, 1993). The few published records of *Neffapis* are from the type locality, 6 km S. Vicuña in southern Elqui Province, and from Las Breas in northern Limarí Province (Rozen and Ruz, 1995). Additional collections at the AMNH are as follows: Chile: Elqui Province: 8 males, 7 km S of Vicuña, X-6-1997 (J. G. Rozen, Jr., H. Navarrete); 2 females, 3 males, same except on *Malesherbia humilis*; 1 male, 6 km S of Vicuña, XI-20-1991 (J. G. Rozen, Jr., L. E. Peña, A. Ugarte); 7 females, 2 males, same except X-30-2000 (J. G. Rozen, Jr.); 1 male, 22 km S of Vicuña, 30°; 10°27'S 70°30'54"W, X-31-2000 (J. G. Rozen, Jr.).

COMMENTS: This group was recognized as one of nine subtribes of an expanded tribe Panurgini by Ascher (2004), along with *Protomelitturgina* (herein considered a tribe, *sensu* Engel), *Melitturgulina*, *Melitturgina*, *Mermiglossina* (herein considered a junior subjective synonym of *Melitturgina*, *sensu* Engel), *Camptopoeina*, *Panurginina*, *Panurgina*, and *Perditina*. The other three panurgine tribes recognized by Ascher (2004) were *Nolanomelissini*, *Calliopsini*, and *Protandrenini*.

Neffapis is apparently an oligolege of *Malesherbia humilis* Poeppig (Malesherbiaceae) (Rozen and Ruz, 1995) and the long tongue of *Neffapis* is probably an adaptation for collecting nectar from this plant. *Neffapis* flies in the austral spring; dates of collection range from 6 October–5 January.

AFRODASYPODINI Engel, new tribe

TYPE GENUS: *Afrodasyopoda* Engel, new genus.

DIAGNOSIS: Glossal apex attenuate; labrum less than four times as long as wide. F1 approximately 2.5 times as long as wide. Compound eyes diverging below. Metatibia lacking keirotichiae; primitively possesses projection on metabasitarsus (like *Promelittini* and some *Sambini*). Basal tergal bands only present on metasomal T2 and T3; median el-

evated area of pygidial plate absent (present in *Dasypodaini*).

COMMENTS: The single species, known only from South Africa, presently known of *Afrodasypodini* is interesting in its intermingling of some traits of the tribes *Dasypodaini* and *Sambini*.

Afrodasyopoda Engel, new genus

TYPE SPECIES: *Rhinochaetula plumipes* Friese, 1912.

DIAGNOSIS: As for the tribe (*vide supra*).

ETYMOLOGY: The new genus-group name is a combination of Africa and the generic name *Dasypoda*, type genus for the subfamily. The name is feminine.

COMMENTS: The only included species is *Afrodasyopoda plumipes* (Friese), **new combination**.

HESPERAPINA Ascher and Engel,
new subtribe

TYPE GENUS: *Hesperapis* Cockerell, 1898.

DIAGNOSIS: Galeal comb present, albeit weak in the American clade of *Hesperapis* and apomorphically reduced to about five bristles in a species of *Eremaphanta* (*Popovapis*). Scopa largely confined to outer surface of metatibia and metabasitarsus; inner metatibial surface with longitudinal median band of keirotichia (scopal setae dense, long, minutely barbed, present on both inner and outer surfaces of metatibia and metabasitarsus, and keirotichia absent in *Dasypoda*); metabasitibial plate present in female and nearly all males (apomorphically absent in some male *Hesperapis*) (absent in *Dasypodaina*). Propodeal profile nearly horizontal at base (all more or less in same plane in *Dasypoda*). Male S1 with broad, transparent marginal zone with deep median cleft; lateroapical lobes of male S7 absent (present in other *dasypodaines*). Gonostylus usually short and broad, fully fused to gonocoxite (nearly always deeply bifid and well differentiated from gonocoxite by narrow, partly membranous area in *Dasypodaina*). Larva with extreme reduction of cephalic and mouthpart structures; maxilla and labium fused; prementum and postmentum fused (Rozen and McGinley, 1974; Rozen, 1978; however, larvae of *Eremaphanta* remain un-

discovered and so validity of larval traits await confirmation).

COMMENTS: *Hesperapina* is herein proposed for *Hesperapis* s.l. (i.e., *sensu* Michener, 2000, including subgenera *Capicola* and *Xeralictoides*; previously these taxa had been treated as separate genera, e.g., Michener, 1981) and *Eremaphanta*. The distribution of hesperapines is unique among bees and features remarkable disjunctions. Nearly all species occur in or near xeric or Mediterranean-climate areas. In the Old World *Eremaphanta* is restricted to xeric regions of Central Asia southwest to Iran (subgenus *Popovapis* to Baluchistan), while *Hesperapis* (*Capicola*) and its sister subgenus *Capicoloides* are restricted to xeric parts of South Africa and Namibia. The remaining subgenera of *Hesperapis* are restricted to North America. Most species are found in seasonally dry habitats in the western United States and northwestern Mexico; the overall range extends from Oregon, North Dakota, and Illinois south to the Gulf Coast of Alabama and northwestern Florida and to Baja California Sur and Morelos, Mexico (Michener, 2000). The only plausible explanation for the extreme disjunction between the three areas of hesperapine occurrence, which are not linked directly by geology, is extensive extinction of ancestral populations in intervening areas due to climatic deterioration (e.g., Engel, 2001). *Hesperapina* thus exhibits limited, relict distribution in three of Sclater's biogeographic regions, whereas its sister group *Dasypodaina* is widely distributed across the Palearctic Region but absent from North America and sub-Saharan Africa.

MACROGALEINA Engel, new subtribe

TYPE GENUS: *Macrogalea* Cockerell, 1930b.

DIAGNOSIS: Body robust and densely setose. Male compound eyes enlarged. Jugal lobe of hind wing greatly enlarged. Female with fasciae of appressed setae on T2–5. Body of mature larva with numerous short setae (some setae hooked at apices), lacking tubercles and elongate setae of subtribe *Allopadina*.

COMMENTS: Species of *Macrogalea* are unlike those of any other *allopapine* genera and

the segregation of the genus into its own subtribe highlights the structural as well as biological differences between it and other alodapines. This distinction is also supported by molecular studies (e.g., Schwarz et al., 2003). The subtribe occurs in Madagascar and Africa (Ethiopia south to Namibia).

ANCYLOSCELIDINA Engel and Michener,
new subtribe

TYPE GENUS: *Ancylloscelis* Latreille, 1829.

DIAGNOSIS: Clypeus strongly protuberant; paraocular carina present along inner margin of compound eye; maxillary palpus with sparse, short setae; scopa on metatibia and metabasitarsus composed of elongate, coarsely plumose setae; cu-a in hind wing less than half as long as second abscissa M + Cu; second abscissa M + Cu in hind wing three-fourths as long as M; male T7 apically rounded; male hind leg greatly modified, metafemur dilated, at least twice as wide as mesofemur; male S7 with broad disc, with 2–4 small apical lobes, shorter than disc.

COMMENTS: This subtribe has been recognized in earlier studies of apine classification but the family-group name has not previously been made available. This minor nomenclatorial difficulty is corrected here.

ACKNOWLEDGMENTS

I am grateful to Michael Oehl, Antonio Arillo, and John S. Ascher for critical reviews of the manuscript. Partial support for this study was provided by NSF EF-0341724. This is contribution Nr. 3408 of the Division of Entomology, Natural History Museum, University of Kansas.

REFERENCES

- Alexander, B.A. 1990. A cladistic analysis of the nomadine bees (Hymenoptera: Apoidea). *Systematic Entomology* 15: 121–152.
- Alexander, B.A., and C.D. Michener. 1995. Phylogenetic studies of the families of short-tongued bees (Hymenoptera: Apoidea). *University of Kansas Science Bulletin* 55(11): 377–424. [publ. date: 30 September 1995]
- André, E. 1886. *Species des Hyménoptères d'Europe & d'Algérie* [vol. 3, fasc. 24–26]: 1–104. Beaune (Côte-d'Or): Chez l'auteur.
- Antropov, A.V. 2000. Digger wasps (Hymenoptera, Sphecidae) in Burmese amber. *Bulletin of the Natural History Museum, London (Geology)* 56(1): 59–77.
- Ascher, J.S. 2003. Appendix: Evidence for the phylogenetic position of *Nolanomelissa* from nuclear EF-1 α sequence data. In G.A.R. Melo and I. Alves-dos-Santos (editors), *Apoidea Neotropica: Homenagem aos 90 Anos de Jesus Santiago Moure*: 107–108. Criciúma: Editora UNESC [Universidade do Extremo Sul Catarinense], xvi + 320 pp.
- Ascher, J.S. 2004. Systematics of the bee family Andrenidae (Hymenoptera: Apoidea). Ph.D. diss., Cornell University, Ithaca, NY, ix + 332 pp.
- Ashmead, W.H. 1890. On the Hymenoptera of Colorado; descriptions of new species, notes, and a list of the species found in the state. *Bulletin of the Colorado Biological Association* 1: 1–47.
- Ashmead, W.H. 1898. Some new genera of bees. *Psyche* 8: 282–285.
- Ashmead, W.H. 1899a. Classification of the bees, or the superfamily Apoidea. *Transactions of the American Entomological Society* 26: 49–100.
- Ashmead, W.H. 1899b. Classification of the entomophilous wasps, or the superfamily Sphegoidea. *Canadian Entomologist* 31: 145–155, 161–174, 212–225, 238–251, 291–300, 322–330, 345–357.
- Baker, D.B. 1994. The dates of the Hymenoptera sections of Costa's *Fauna del Regno di Napoli*. *Archives of Natural History* 21(3): 351–356.
- Beebe, W. 1925. Studies of a tropical jungle; one quarter of a square mile of jungle at Kartabo, British Guiana. *Zoologica* 6: 5–193.
- Berthold, A.A. 1827. Latreille's natürliche Familien der Tierreichs, aus dem Französischen mit Anmerkungen und Zusätzen. Weimar: Industr.-Compt., viii + 602 pp.
- Bohart, G.E. 1951. Tribe Tachytini. In C.F.W. Muesebeck, K.V. Krombein, and H.K. Townes (editors), *Hymenoptera of American North of Mexico—Synoptic Catalog*: 945–953. Washington: United States Department of Agriculture [Agriculture Monograph 2], 1420 pp.
- Bohart, R.M. 1966. A review of *Aphilanthops* and related genera (Hymenoptera: Sphecidae). *Proceedings of the Entomological Society of Washington* 68: 158–167.
- Bohart, R.M., and D.R. Horning. 1971. California bembicine sand wasps. *Bulletin of the California Insect Survey* 13: 1–49.
- Bohart, R.M., and A.S. Menke. 1963. A reclassification of the Sphecinae, with a revision of the Nearctic species of the tribes Sceliphronini and Sphecini (Hymenoptera, Sphecidae). Uni-

- versity of California Publications in Entomology 30(2): 91–182.
- Bohart, R.M., and A.S. Menke. 1976. Sphecoid wasps of the world: a generic revision. Berkeley: University of California, ix + 695 pp.
- Börner, C. 1919. Stammesgeschichte der Hautflügler. Biologisches Zentralblatt 39(4): 145–186.
- Bour, R., and A. Dubois. 1984. Nomenclature ordinaire et familiale des tortues (Reptilia). Studia Geologica Salmanticensia (Studia Palaeoethnologia I) 1(especial): 77–86.
- Bradley, J.C. 1926. Hymenoptera. In M.D. Leonard (editor), A list of the insects of New York with a list of spiders and certain other allied groups: 870–1033. Memoirs of the Cornell University Agriculture Experiment Station 101: 1–1121.
- Brady, S.G., and B.N. Danforth. 2004. Recent intron gain in elongation factor-1 α of colletid bees (Hymenoptera: Colletidae). Molecular Biology and Evolution 21(4): 691–696.
- Brauns, H. 1902. *Eucondylops* n. g. apidarum. Zeitschrift für Systematische Hymenopterologie und Dipterologie 2: 377–380.
- Brèthes, J. 1909. Hymenoptera paraguayensis. Anales del Museo Nacional de Buenos Aires 20: 225–256.
- Brothers, D.J. 1975. Phylogeny and classification of the aculeate Hymenoptera, with special reference to Mutillidae. University of Kansas Science Bulletin 50(11): 483–648. [publ. date: 15 August 1975]
- Brues, C.T., and A.L. Melander. 1932. Classification of insects. Bulletin of the Museum of Comparative Zoology 73: 1–672.
- Burmeister, C.H.C. 1829. De Insectorum Systemate Naturali. Halle: Gruneret, [iv] + 40 + [3] pp.
- Cameron, P. 1903. Descriptions of new species of Hymenoptera taken by Mr. Edward Whymper on the “Higher Andes of the Equator”. Transactions of the American Entomological Society 29: 225–238.
- Cockerell, T.D.A. 1896. The bees of the genus *Andrena* found in New Mexico. Annals and Magazine of Natural History, series 6, 18: 78–92.
- Cockerell, T.D.A. 1897. On the Mexican bees of the genus *Augochlora*. Canadian Entomologist 29: 4–6.
- Cockerell, T.D.A. 1898. New bees from New Mexico. Canadian Entomologist 30: 146–148.
- Cockerell, T.D.A. 1900a. Observations on bees collected in Las Vegas, New Mexico, and in the adjacent mountains. Annals and Magazine of Natural History, series 7, 5: 401–416.
- Cockerell, T.D.A. 1900b. Descriptions of new bees collected by Mr. H. H. Smith in Brazil—I. Proceedings of the Academy of Natural Sciences of Philadelphia 52: 356–377.
- Cockerell, T.D.A. 1902. Appendix. In T.D.A. Cockerell and E. Atkins, Contributions from the New Mexico Biological Station—XII. On some genera of bees: 233–234. Annals and Magazine of Natural History, series 7, 9: 230–234.
- Cockerell, T.D.A. 1905. Descriptions and records of bees—III. Annals and Magazine of Natural History, series 7, 16: 301–308.
- Cockerell, T.D.A. 1906. The North American bees of the family Anthophoridae. Transactions of the American Entomological Society 32: 63–116.
- Cockerell, T.D.A. 1907. Descriptions and records of bees—XV. Annals and Magazine of Natural History, series 7, 20: 59–68.
- Cockerell, T.D.A. 1908a. Notes on the bee-genus *Exaerete*. Psyche 15: 41–42.
- Cockerell, T.D.A. 1908b. Descriptions and records of bees—XX. Annals and Magazine of Natural History, series 8, 2: 323–334.
- Cockerell, T.D.A. 1909a. Some European fossil bees. Entomologist 42: 313–317.
- Cockerell, T.D.A. 1909b. Descriptions of Hymenoptera from Baltic amber. Schriften der Physikalisch-ökonomischen Gesellschaft, Königsberg 50(1): 1–20.
- Cockerell, T.D.A. 1909c. Descriptions and records of bees—XXIII. Annals and Magazine of Natural History, series 8, 4: 393–404.
- Cockerell, T.D.A. 1926. Descriptions and records of bees—CVIII. Annals and Magazine of Natural History, series 9, 17: 214–226.
- Cockerell, T.D.A. 1930a. The bees of Australia. Australian Zoologist 6(2): 137–156.
- Cockerell, T.D.A. 1930b. A new African genus of Ceratinidae (Hymenoptera, Apoidea). Revue de Zoologie et de Botanique Africaines 18: 291–293.
- Cockerell, T.D.A. 1932. Descriptions and records of bees—CXXXI. Annals and Magazine of Natural History, series 10, 9: 355–365.
- Cockerell, T.D.A. 1934. The bees of Australia [continued]. Australian Zoologist 8(1): 1–38.
- Cockerell, T.D.A., and E. Atkins. 1902. Contributions from the New Mexico Biological Station—XIII. On the bees of the family Nomadidae of Ashmead. Annals and Magazine of Natural History, series 7, 10: 40–46.
- Cockerell, T.D.A., and W. P. Cockerell. 1901. Contributions from the New Mexico Biological Station—IX. On certain genera of bees. Annals and Magazine of Natural History, series 7, 7: 46–50.
- Cockerell, T.D.A., and W.W. Robbins. 1910. An

- introduction to the study of Rocky Mountain bees. University of Colorado Studies 7: 179–195.
- Costa, A. 1858. Immenotteri aculeati, familia degli Sfecidei. In O.G. Costa and A. Costa, Fauna del Regno di Napoli ossia enumerazione di tutti gli animali che abitano le diverse regioni di questo regno e le acque che le bagnano e descrizione de'nuovi o poco esattamente conosciuti con figure ricavate da originali viventi e dipinte al naturale [Parts I & II]: 1–28. Napoli [Naples]: Antonio Cons, 36 + 36 + 39 + 69 + 56 + 38 + 16 + 8 + 6 pp., 27 pls. [publ. date: *vide* Baker (1994)]
- Costa, A. 1859. Immenotteri aculeati, familia de Nissonidei; Nyssonidea (Nyssonidae, Dahlb.). In O.G. Costa and A. Costa, Fauna del Regno di Napoli ossia enumerazione di tutti gli animali che abitano le diverse regioni di questo regno e le acque che le bagnano e descrizione de'nuovi o poco esattamente conosciuti con figure ricavate da originali viventi e dipinte al naturale [Parts I & II]: 1–56. Napoli [Naples]: Antonio Cons, 36 + 36 + 39 + 69 + 56 + 38 + 16 + 8 + 6 pp., 27 pls. [publ. date: *vide* Baker (1994)]
- Crawford, J.C. 1916. Some new American Hymenoptera. Insecutor Inscitiae Menstruus 4: 135–144.
- Cresson, E.T. 1887. Synopsis of the families and genera of the Hymenoptera of America, north of Mexico, together with a catalogue of the described species, and bibliography. Transactions of the American Entomological Society, supplement 1887: viii + 1–350 + [1: errata].
- Curtis, J. 1833. British entomology: being illustrations and descriptions of the genera of insects found in Great Britain and Ireland [vol. 10]. London: Privately printed, pl. 434–481.
- Dahlbom, A.G. 1835. Clavis Novi Hymenopterorum Systematis adjecta Synopsi Larvarum ejusdem Ordinis Scandinavicarum Eruciformium. Lund: Berling, v + 40 pp., 1 pl.
- Dahlbom, A.G. 1842. Dispositio methodica specierum Scandinavicarum ad familias Hymenopterorum naturales pertinentium [part 1]. *Sphex* in sensu Linnaeano. Lund: Berling, 16 pp.
- Dalla Torre, C.G., de [K. W., von]. 1896. Catalogus Hymenopterorum hucusque descriptorum systematicus et synonymicus. Volumen X: Apidae (Anthophila). Lipsiae [Leipzig]: Engelmann, viii + 643 pp.
- Dalla Torre, C.G., de [K. W., von]. 1897. Catalogus Hymenopterorum hucusque descriptorum systematicus et synonymicus. Volumen VIII: Fossores. Lipsiae [Leipzig]: Engelmann, viii + 749 pp. [publ. date: no later than October 1897]
- Dalla Torre, C.G., de [K. W., von], and H. Friese. 1894. Synonymischer Katalog der europäischen Schmarotzerbienen. Entomologische Nachrichten 20: 33–43.
- Danforth, B.N., S.G. Brady, S.D. Sipes, and A. Pearson. 2004. Single-copy nuclear genes recover Cretaceous-age divergences in bees. Systematic Biology 53(2): 309–326.
- Ducke, A. 1902. Ein neues subgenus von *Halictus* Latr. Zeitschrift für Systematische Hymenopterologie und Dipterologie 2: 102–103.
- Ducke, A. 1912. Die natürlichen Bienengenera Südamerikas. Zoologische Jahrbücher, Abteilung für Systematik, Geographie und Biologie der Tiere 34: 51–116.
- Dupuis, C. 1986. Dates de publication de l'“Histoire naturelle générale et particulière des Crustacés et des insectes” (1802–1805) par Latreille dans le “Buffon de Sonnini”. Annales de la Société de Entomologique de France 22: 205–210.
- Eardley, C.D. 1996. The genus *Scapter* Lepeletier & Serville (Hymenoptera: Colletidae). African Entomology 4(1): 37–92.
- Engel, M.S. 1997. *Ischnomelissa*, a new augochlorine bee genus (Halictidae) from Colombia. Studies on Neotropical Fauna and Environment 32(1): 41–46. [publ. date: November 1997]
- Engel, M.S. 1998a. A new species of the Baltic amber bee genus *Electrapis* (Hymenoptera: Apidae). Journal of Hymenoptera Research 7(1): 94–101. [publ. date: 14 May 1998]
- Engel, M.S. 1998b. Comparative morphology and the phylogeny of the corbiculate bees (Hymenoptera: Apidae; Apinae). In M.P. Schwarz and K. Hogendoorn (editors), Social insects at the turn of the millennium: Proceedings of the XIII International Congress of IUSSI [International Union for the Study of Social Insects]: 151. Adelaide: Flinders University Press, [v] + 535 pp. [publ. date: 29 December 1998]
- Engel, M.S. 1999a. Augochlorini Beebe, 1925 (Insecta, Hymenoptera): Corrected authorship and date (not Moure, 1943). Bulletin of Zoological Nomenclature 56(3): 198.
- Engel, M.S. 1999b. *Megachile glaesaria*, the first megachilid bee fossil from amber (Hymenoptera: Megachilidae). American Museum Novitates 3276: 1–13. [publ. date: 12 November 1999]
- Engel, M.S. 2000. Classification of the bee tribe Augochlorini (Hymenoptera: Halictidae). Bulletin of the American Museum of Natural History 250: 1–88. [publ. date: 7 April 2000]
- Engel, M.S. 2001. A monograph of the Baltic amber bees and evolution of the Apoidea (Hymenoptera). Bulletin of the American Museum

- of Natural History 259: 1–192. [publ. date: 16 April 2001]
- Engel, M.S. 2004. Fideiine phylogeny and classification revisited (Hymenoptera: Megachilidae). *Journal of the Kansas Entomological Society* 77(4): 821–836. [publ. date: 10 December 2004]
- Engel, M.S., and K. Krishna. 2004. Family-group names for termites (Isoptera). *American Museum Novitates* 3432: 1–9. [publ. date: 27 February 2004]
- Evans, H.E. 1959. Studies on the larvae of digger wasps, part V: Conclusion. *Transactions of the American Entomological Society* 85: 137–191.
- Fabricius, J.C. 1775. *Systema Entomologiae, sistens insectorum classes, ordines, genera, species, adiectis synonymis, locis, descriptionibus, observationibus*. Flensburgi et Lipsiae [Flensburg and Leipzig]: Korte, xxxii + 832 pp.
- Fabricius, J.C. 1793. *Entomologia Systematica emendata et aucta: Secundum classes, ordines, genera, species adiectis synonymis, locis, observationibus, descriptionibus* [vol. 2]. Hafniae [Copenhagen]: Proft, viii + 519 pp.
- Fabricius, J.C. 1804. *Systema Piezatorum Secundum Ordines, Genera, Species, adiectis Synonymis, Locis, Observationibus, Descriptionibus*. Brunswickae [Brunswick]: Reichard, 439 + 30 pp.
- Fallén, C.F. 1813. *Specimen Novam Hymenoptera Disponendi Methodum Exhibens*. Lund: Berling, 42 pp., 1 pl.
- Fernald, H.T. 1905. The type of the genus *Sphex*. *Entomological News* 16: 163–166.
- Fox, W.J. 1894 [1895]. A proposed classification of the fossorial Hymenoptera of North America. *Proceedings of the Academy of Natural Sciences of Philadelphia* 1894(3): 292–307. [publ. date: January 1895]
- Friese, H. 1898. Beiträge zur Bienenfauna von Aegypten. *Természetrajzi Füzetek* 21: 303–313.
- Friese, H. 1899. Monographie der Bienengattungen *Argogorytina*, *Caupolicana* und *Oxaea*. *Annalen des Kaiserlich-königlichen Naturhistorischen Hofmuseum (Wien)* 14: 239–246.
- Friese, H. 1903a. *Meliturgula*, eine neue Bienen-gattung aus Süd-Afrika (Hym.). *Zeitschrift für Systematische Hymenopterologie und Dipterologie* 3: 33–34.
- Friese, H. 1903b. Neue Meliponiden II (Hym.). *Zeitschrift für Systematische Hymenopterologie und Dipterologie* 3: 359–361.
- Friese, H. 1905. Die Wollbienen Afrikas. Genus *Anthidium*. *Zeitschrift für Systematische Hymenopterologie und Dipterologie* 5: 65–75.
- Friese, H. 1908. Neue Bienenarten aus Ostafrika. *Deutsche Entomologische Zeitschrift* 1908: 567–572.
- Friese, H. 1909. Die Bienen Afrikas nach dem Stande unserer heutigen Kenntnis. In L. Schultze (editor), *Zoologische und anthropologische Ergebnisse einer Forschungsreise im westlichen und zentralen Südafrika ausgeführt in den Jahren 1903–1905 mit Unterstützung der Kgl. [Königlichen] Preussischen Akademie der Wissenschaften zu Berlin* [Band 2, Lieferung 1, X Insecta (3^{te} Serie): Jenaische Denkschriften 14]: 83–476, pls. ix–x. Jena: Gustav Fischer.
- Friese, H. 1911. Neue Bienen der äthiopischen Region. *Zoologische Jahrbücher, Abteilung für Systematik, Geographie und Biologie der Tiere* 30: 671–678.
- Friese, H. 1912. Neue und wenig bekannte Bienen Süd-Afrikas (Hym.). *Archiv für Naturgeschichte, Abteilung A* 78(5): 181–189.
- Friese, H. 1925. Neue neotropischen Bienenarten, zugleich II. Nachtrag zur Bienenfauna von Costa Rica (Hym.). *Stettiner Entomologische Zeitung* 86: 1–41.
- Frison, T.H. 1919. Keys for the separation of the Bremidae, or bumblebees, of Illinois, and other notes. *Transactions of the Illinois State Academy of Science* 12: 157–165.
- Gerstäcker, A. 1857. [Bees and wasps collected in Mozambique]. *Monatsberichte, Akademie der Wissenschaften, Berlin* 1857(29 October): 460–464.
- Giraud, J. 1863. Hyménoptères recueillis aus environs de Suse, en Piémont, et dans le département des Hautes-Alpes, en France; et description de quinze espèces nouvelles. *Verhandlungen der Zoologisch-Botanischen Gesellschaft in Wien* 13: 11–46.
- Gittins, A.R. 1969. Revision of the Nearctic Psenini (Hymenoptera: Sphecidae). I. Redescriptions and keys to the genera and subgenera. *Transactions of the American Entomological Society* 95: 49–76.
- Guérin-Méneville, F.E. 1844. *Iconographie du Règne Animal de G. Cuvier; ou, représentation d'après nature de l'une des espèces les plus remarquables, et souvent non encore figurées, de chaque genre d'animaux: Avec un texte descriptif mis au courant de la science: Ouvrage pouvant servir d'atlas à tous les traités de zoologie* [tome 3: Texte explicatif]. Paris: Baillière, 576 pp., 106 pls.
- Handlirsch, A. 1925. Systematische Übersicht. In C. Schröder (editor), *Handbuch der Entomologie*, Band 3: Geschichte, Literatur, Technik, Paläontologie, Phylogenie, Systematik: 377–1143. Jena: Fischer, viii + 1201 pp.
- von Hoffmannsegg, C.J.C. 1817. *Entomologische Bemerkungen bei Gelegenheit der Abhandlungen über amerikanische Insekten*. *Zoologisches Magazin, Wiedemann* [Kiel] 1: 8–56.

- Holmberg, E.L. 1886. Sobre ápidos nómadas de la República Argentina. *Anales de la Sociedad Científica Argentina* 22: 231–240, 272–286.
- Holmberg, E.L. 1903. *Delectus Hymenopterologicus Argentinus: Hymenopterorum Argentinorum et quorundam exoticorum observationes synonymicas, addendas, novorumque generum specierumque descriptiones continens*. *Anales del Museo Nacional de Buenos Aires* 2: 377–512.
- Illiger, K. 1801. Namen der Insekten-Gattungen, ihr Genitiv, ihr grammatisches Geschlecht, ihr Silbenmass, ihre Herleitung: Zugleich mit den Deutschen Benennungen. *Magazin für Insektenkunde* 1: 125–155.
- Illiger, K. 1806. William Kirby's Familien der Bienenartigen Insekten mit Zusätzen, Nachweisungen und Bemerkungen. *Magazin für Insektenkunde* 5: 28–175.
- International Commission on Zoological Nomenclature. 1939. Opinion 135. The suppression of the so-called "Erlangen List" of 1801. *Opinions Rendered by the International Commission on Zoological Nomenclature* 2: 9–12.
- International Commission on Zoological Nomenclature. 1974. Opinion 1005. *Trypeta* Meigen, 1803 (Insecta: Diptera); *Trypetes* Schoenherr, 1836 (Insecta: Coleoptera); *Trypetesa* Norman, 1903 (Crustacea: Cirripedia): Removal of homonymy in family-group names under the plenary powers. *Bulletin of Zoological Nomenclature* 30: 151–152.
- International Commission on Zoological Nomenclature. 1979. Opinion 1115. Validation of the generic name *Nysson* Latreille (Hymenoptera, Sphecidae) as from 1796. *Bulletin of Zoological Nomenclature* 35: 175–179.
- International Commission on Zoological Nomenclature. 1985. International Code of Zoological Nomenclature [Third Edition]. London: International Trust for Zoological Nomenclature, xx + 338 pp.
- International Commission on Zoological Nomenclature. 1987. Opinion 1445. Heterogynidae Rambur, 1866 (Insecta, Lepidoptera) and Heterogynidae Nagy, 1969 (Insecta, Hymenoptera): A ruling to remove the homonymy. *Bulletin of Zoological Nomenclature* 44: 150–151.
- International Commission on Zoological Nomenclature. 1993. Opinion 1713. Some bee family-group names (Insecta, Hymenoptera: Names based on *Colletes* Latreille, 1802, on *Paracolletes* Smith, 1853, on *Halictus* Latreille, 1804, on *Anthidium* Fabricius, 1804 and on *Anthophora* Latreille, 1803 given precedence over some senior names. *Bulletin of Zoological Nomenclature* 50(1): 85–89.
- International Commission on Zoological Nomenclature. 1999a. International Code of Zoological Nomenclature [Fourth Edition]. London: International Trust for Zoological Nomenclature, xxix + 306 pp.
- International Commission on Zoological Nomenclature. 1999b. Opinion 1926. Dasypodidae Börner, 1919 (Insecta, Hymenoptera): Spelling emended to Dasypodaidae, so removing the homonymy Dasypodidae Gray, 1821 (Mammalia, Xenarthra). *Bulletin of Zoological Nomenclature* 56(2): 156–157.
- International Commission on Zoological Nomenclature. 2000. Opinion 1952. Augochlorini Beebe, 1925 (Insecta, Hymenoptera): Given precedence over Oxystoglossini Schrottky, 1909. *Bulletin of Zoological Nomenclature* 57(2): 128–129.
- Jörgensen, P. 1909. Beobachtungen über blumenbesuch, Biologie, Verbreitung usw. der Bienen von Mendoza (Hym.). *Deutsche Entomologische Zeitschrift* 1909: 53–65, 211–228.
- Jurine, L. 1807. *Nouvelle Méthode de Classer les Hyménoptères et les Diptères*. Geneva: Paschoud, 320 + 4 pp., 14 pls.
- Kirby, W. 1802. *Monographia Apum Angliæ; or, an attempt to divide into their natural genera and families, such species of the Linnean genus Apis as have been discovered in England: with descriptions and observations*. Ipswich: White, vol. 1: xxii + 258 pp., pls. 1–14; vol. 2: [i] + 388 pp., pl. 15–18.
- Kirby, W. 1837. Part IV, Insects. In J. Richardson (editor), *Fauna Boreali-Americana, or, the zoology of the northern parts of British America; containing descriptions of the objects of natural history collected on the late northern land expeditions, under command of Captain Sir John Franklin, R.N.*: 1–325. London: Longman, xxxix + [i] + 325 pp., 8 pls.
- Kirby, W., and W. Spence. 1826. *An introduction to entomology, or elements of the natural history of insects* [vol. 3]. London: Longman, v + [iii] + 732 pp.
- Klug, J.C.F. 1807a. *Oxaea*, eine neue Gattung aus der Ordnung der Piezaten. *Gesellschaft Naturforschender Freunde zu Berlin, Magazin für neuesten Entdeckungen in der gesammten Naturkunde* 1: 261–263.
- Klug, J.C.F. 1807b. Kritische Revision der BienenGattungen in Fabricius neuem Piezaten-systeme. *Magazin für Insektenkunde* 6: 200–228.
- Klug, J.C.F. 1810. Einige neue PiezatenGattungen. *Gesellschaft Naturforschender Freunde zu Berlin, Magazin für neuesten Entdeckungen in der gesammten Naturkunde* 4: 31–45, pl. 1.
- Lacordaire, J.T. 1866. *Histoire Naturelle des Insectes. Genera des Coleopteres* [vol. 7]. Paris: Roret, 620 pp.

- Latreille, P.A. 1802a. Histoire Naturelle des Fourmis, et recueil de memoires et d'observations sur les abeilles, les araignées, les faucheurs, et autres insectes. Paris: Crapelet, xvi + 445 pp. [publ. date: April 1802]
- Latreille, P.A. 1802b. Histoire Naturelle, Générale et Particulière des Crustacés et des Insectes. Ouvrage faisant suite à l'histoire naturelle générale et particulière, composée par Leclerc de Buffon, et rédigée par C. S. Sonnini, membre de plusieurs sociétés savantes. Tome troisième [3]. Paris: Dufart, xii + 467 pp. [publ. date: November 1802; despite reading "X" on the title page, this volume was published in XI of the "metric" Revolutionary Calendar which lasted from 22 September 1802–21 September 1803; *vide* Dupuis (1986)]
- Latreille, P.A. 1803. [Various sections]. In Nouveau Dictionnaire d'Histoire Naturelle, appliquée aux arts, principalement à l'agriculture et à l'économie rurale et domestique, tome 18 [vol. 18]. Paris: Déterville.
- Latreille, P.A. 1804. Tableau méthodique des insectes. In Nouveau Dictionnaire d'Histoire Naturelle, appliquée aux arts, principalement à l'agriculture et à l'économie rurale et domestique, tome 24 [vol. 24], caractères et tables: 129–200. Paris: Déterville, 84 + 85 + 238 + 18 + 34 pp.
- Latreille, P.A. 1805. Histoire Naturelle, Générale et Particulière des Crustacés et des Insectes. Ouvrage faisant suite à l'histoire naturelle générale et particulière, composée par Leclerc de Buffon, et rédigée par C. S. Sonnini, membre de plusieurs sociétés savantes. Tome treizième [13]. Paris: Dufart, 432 pp., pls. 98–103. [publ. date: 20 February–21 March 1805 = "An XIII" of the Revolutionary Calendar]
- Latreille, P.A. 1807. Genera crustaceorum et insectorum secundum ordinem naturalem in familias disposita, iconibus exemplisque plurimis explicata [vol. 3]. Paris: Koenig, 258 pp.
- Latreille, P.A. 1809. Genera crustaceorum et insectorum secundum ordinem naturalem in familias disposita, iconibus exemplisque plurimis explicata, tome 4. Paris: Koenig, 399 pp.
- Latreille, P.A. 1810. Considérations générales sur l'ordre naturel des animaux composant les classes des crustacés, des arachnides, et des insectes; avec un tableau méthodique de leurs genres, disposés en familles. Paris: Schoell, 444 pp.
- Latreille, P.A. 1829. Les Insectes. In G.C.L.D. Cuvier (editor), Le règne animal distribué d'après son organisation pour servir de base à l'histoire naturelle des animaux et à l'introduction à l'anatomie comparée [2nd Edition, Vol. 5]. Paris: Déterville, xxiv + 556 pp.
- Leach, W.E. 1815. Entomology. In D. Brewster (editor), The Edinburgh Encyclopaedia [Vol. 9, part 1]: 57–172. Edinburgh: Blackwood, [2] + 766 pp.
- Lepeletier de Saint Fargeau, A.L.M. 1832 [1833]. Observations sur l'ouvrage intitulé: "Bombi Scandinaviae Monographice Tractati, etc., à Gustav Dahlbom. Londini Gothorum, 1832;" auxquelles on a joint les caractères des genres *Bombus* et *Psithyrus*, et la description des especes qui appartiennent au dernier. Annales de la Société Entomologique de France 1: 366–382.
- Lepeletier de Saint Fargeau, A.L.M. 1836. Histoire Naturelle des Insectes—Hyménoptères [vol. 1]. Paris: Roret, 547 pp.
- Lepeletier de Saint Fargeau, A.L.M. 1841. Histoire Naturelle des Insectes—Hyménoptères [vol. 2]. Paris: Roret, 680 pp.
- Lepeletier de Saint Fargeau, A.L.M. 1845. Histoire Naturelle des Insectes—Hyménoptères [vol. 3]. Paris: Roret, 644 pp.
- Lepeletier de Saint Fargeau, A.L.M., and J.G. Audinet-Serville. 1825. [Sections]. In G.A. Olivier (editor), Encyclopédie Méthodique, ou par ordre de matières. Histoire Naturelle. Insectes [vol. 10, part 1]. Paris: Agasse, 1–344 pp.
- Lepeletier de Saint Fargeau, A.L.M., and J.G. Audinet-Serville. 1828. [Sections]. In G.A. Olivier (editor), Encyclopédie Méthodique, ou par ordre de matières. Histoire Naturelle. Insectes [vol. 10, part 2]. Paris: Agasse, 345–832 pp.
- Linnaeus, C. 1758. Systema Naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis synonymis, locis [10th Edition, revised]. Holmiae [Stockholm]: Laurentii Salvii, 824 pp.
- Linsley, E.G., and C.D. Michener. 1937. Some new genera and species of North American parasitic bees (Hymenoptera). Pan-Pacific Entomologist 13: 75–84.
- Linsley, E.G., and C.D. Michener. 1939. A generic revision of the North American Nomadidae (Hymenoptera). Transactions of the American Entomological Society 65: 265–305. [Publ. date: 24 October 1939]
- Loew, H. 1861. Ueber die Dipterenfauna des Bernsteins. Amtlicher Bericht über die fünf und dreissigste Versammlung Deutscher Naturforscher und Ärzte in Königsberg in Preussen im September 1860. 1861: 88–98.
- Lomholdt, O. 1985. A reclassification of the larine tribes with a revision of the Miscophini of southern Africa and Madagascar (Hymenoptera: Sphecidae). Entomologica Scandinavica, supplement 24: 1–183.
- Marikovskaya, T.P. 1976. On the systematics of the tribe Anthophorini (Hymenoptera, Apoidea,

- Anthophoridae). *Entomologicheskoe Obozrenie* 55: 684–690. [In Russian]
- McGinley, R.J. 1981. Systematics of the Colletidae based on mature larvae with phenetic analysis of apoid larvae (Hymenoptera: Apoidea). University of California Publications in Entomology 91: 1–307.
- McKenna, M.C., and S.K. Bell. 1997. Classification of mammals above the species level. New York: Columbia University Press, xii + [1] + 631 pp.
- Meigen, J.W. 1803. Versuch einer neuen Gattungs-Eintheilung der europäischen zweiflügeligen Insekten. *Magazin für Insektenkunde* 2: 259–281. [publ. date: after 17 May 1803]
- Melo, G.A.R. 1999. Phylogenetic relationships and classification of the major lineages of Apoidea (Hymenoptera), with emphasis on the crabronid wasps. University of Kansas Museum of Natural History Scientific Papers 14: 1–55.
- Menke, A.S. 1967. *Odontosphex* Arnold, a genus of the Philanthinae, with a key to the tribes and genera of the subfamily (Hymenoptera: Sphecidae). *Pan-Pacific Entomologist* 43: 141–148.
- Menke, A.S. 1968. New South American genera and species of the tribe Bothystethini (Hymenoptera; Sphecidae; Larrinae). *Acta Zoologica Lilloana* 22: 89–99.
- Menke, A.S. 1989. *Arpactophilus* reassessed, with three bizarre new species from New Guinea (Hymenoptera: Sphecidae: Pemphredoninae). *Invertebrate Taxonomy* 2: 737–747.
- Menke, A.S. 1997. Family-group names in Sphecidae (Hymenoptera: Apoidea). *Journal of Hymenoptera Research* 6(2): 243–255. [publ. date: 20 October 1997]
- Menke, A.S., and W. Pulawski. 2002. Authorship of the family-group names Palarini and Xenosphecini (Hymenoptera: Apoidea: Crabronidae). *Journal of Hymenoptera Research* 11: 366–367.
- Michener, C.D. 1941. A generic revision of the American Osmiinae with descriptive notes on Old World genera (Hymenoptera, Megachilidae). *American Midland Naturalist* 26(1): 147–166. [publ. date: July 1941]
- Michener, C.D. 1944. Comparative external morphology, phylogeny, and a classification of the bees (Hymenoptera). *Bulletin of the American Museum of Natural History* 82(6): 151–326. [publ. date: 10 April 1944]
- Michener, C.D. 1948. The generic classification of the anthidiine bees (Hymenoptera, Megachilidae). *American Museum Novitates* 1381: 1–29. [publ. date: 29 July 1948]
- Michener, C.D. 1954. Bees of Panamá. *Bulletin of the American Museum of Natural History* 104(1): 1–175. [publ. date: 21 June 1954]
- Michener, C.D. 1965a. A classification of the bees of the Australian and South Pacific regions. *Bulletin of the American Museum of Natural History* 130: 1–362. [publ. date: 6 December 1965]
- Michener, C.D. 1965b. A generic review of the Dufoureae of the Western Hemisphere (Hymenoptera: Halictidae). *Annals of the Entomological Society of America* 58(3): 321–326.
- Michener, C.D. 1966. The classification of the Diaphaglossinae and North American species of the genus *Caupolicana* (Hymenoptera, Colletidae). *University of Kansas Science Bulletin* 46(20): 717–751. [publ. date: 2 December 1966]
- Michener, C.D. 1981. Classification of the bee family Melittidae, with a review of the species of Meganomiinae. *Contributions of the American Entomological Institute* 18(3): iii + 1–135. [publ. date: 23 December 1981]
- Michener, C.D. 1986a. Family-group names among bees. *Journal of the Kansas Entomological Society* 59(2): 219–234. [publ. date: 28 May 1986]
- Michener, C.D. 1986b. New Peruvian genus and a generic review of Andreninae (Hymenoptera: Apoidea: Andrenidae). *Annals of the Entomological Society of America* 79(1): 62–72.
- Michener, C.D. 1989. Classification of American Colletinae (Hymenoptera, Apoidea). *University of Kansas Science Bulletin* 53(11): 622–703. [publ. date: 25 May 1989]
- Michener, C.D. 1997. Genus-group names of bees and supplemental family-group names. *University of Kansas Museum of Natural History Scientific Papers* 1: 1–81. [publ. date: 25 July 1997]
- Michener, C.D. 2000. *The Bees of the World*. Baltimore: Johns Hopkins University Press, xiv + [1] + 913 pp.
- Michener, C.D., W.E. LaBerge, and J.S. Moure. 1955. Canephorulini, a tribe of South American bees (Hymenopt.-Apoidea). *Dusenian* 6(5): 207–212. [publ. date: 30 September 1955]
- Michener, C.D., and J.S. Moure. 1957. A study of the classification of the more primitive non-parasitic anthophorine bees (Hymenoptera, Apoidea). *Bulletin of the American Museum of Natural History* 112(5): 395–452. [publ. date: 19 August 1957]
- Moure, J.S. 1940. Apoidea neotropica. *Arquivos de Zoologia do Estado de São Paulo* 2: 39–64. [publ. date: 16 November 1940]
- Moure, J.S. 1941. Notas sobre abelhas do grupo *Tetrapedia* Klug (Hym. Apoidea). *Revista de Entomologia* 12(3): 515–521. [publ. date: December 1941]
- Moure, J.S. 1943. Notas sobre abelhas da coleção

- Zikán (Hym. Apoidea). *Revista de Entomologia* 14: 447–484.
- Moure, J.S. 1944. Apoidea da coleção do Conde Amadeu A. Barbiellini (Hym. Apoidea). *Revista de Entomologia* 15(1–2): 1–18.
- Moure, J.S. 1945. Contribuição para o conhecimento dos Diphaglossinae, particularmente *Ptiloglossa* (Hym.–Apoidea). *Arquivos do Museu Paranaense* 4(6): 137–178. [publ. date: April 1945]
- Moure, J.S. 1946. Melíponas do Brasil. *Characas e Quintais* 74: 609–612. [publ. date: 15 November 1946]
- Moure, J.S. 1947. Novos agrupamentos genéricos e algumas espécies novas de abelhas sulamericanas. *Publicações Avulsas Museu Paranaense* 3: 1–37.
- Moure, J.S. 1949. Las especies Chilenas de la sub-familia Lithurginae (Hym.–Apoidea). *Arquivos do Museu Paranaense* 7(5): 265–286. [publ. date: December 1949]
- Moure, J.S. 1950. Euhirbstiinae nova sub-familia de Andrenidae (Hymopt.–Apoidea). *Dusenía* 1(5): 303–306. [publ. date: 30 September 1950]
- Moure, J.S. 1953. Notas sobre Colletidae Sul-Americanos (Hymenoptera–Apoidea). *Dusenía* 4(1): 61–78. [publ. date: 31 January 1953]
- Moure, J.S. 1955. Notas sobre Epeolini sulamericanos (Hymenopt.–Apoidea). *Dusenía* 6(3–4): 115–138. [publ. date: 31 July 1955]
- Moure, J.S. 1992 [1994]. *Lissopedia*, gen. n. de Paratetrapediini para a região neotropical, com as descrições de três espécies novas (Hymenoptera, Apoidea, Anthophoridae). *Revista Brasileira de Zoologia* 9: 305–317. [publ. date: 1994]
- Moure, J.S., and C.D. Michener. 1955. The bee family Fideliidae in South America (Hymenopt.–Apoidea). *Dusenía* 6(5): 199–206. [publ. date: 30 September 1955]
- Nagy, C.G. 1969. A new taxon of the family Heterogynidae Latreille (Hym., Aculeata). *Entomologische Mitteilungen aus dem Zoologischen Staatsinstitut und Zoologischen Museum Hamburg* 64: 299–304.
- Nemokv, P.G., and A.S. Lelej. 1996. Phylogenetic relationships and classification of the digger wasps tribe Gorytini (Hymenoptera: Sphecidae, Nyssoninae). *Far Eastern Entomologist* 37: 1–14.
- Newman, E. 1834. Attempted division of British insects into natural orders. *Entomological Magazine* 2: 379–431.
- Nobile, V., and S. Tomarchio. 2000. Apoidei nuovi o poco noti di alcune regioni d'Italia (Hymenoptera, Apoidea, Megachilidae). I. Bollettino dell' Accademia Gioenia di Scienze Naturali 33(358): 43–54.
- Nylander, W. 1848. Adnotationes in expositionem monographicam apum borealium. *Notiser ur Sällskapetets pro Fauna et Flora Fennica Förhandlingar* 1: 165–282, pl. iii.
- Ogloblin, A.A. 1956. Dos géneros nuevos de la tribu Exomalopsini (Apidae, Hym.) de la República Argentina. *Dusenía* 7(3): 149–158. [publ. date: 30 June 1956]
- Ohl, M. 1996. Die phylogenetischen Beziehungen der Sphecinae (Hymenoptera: Apoidea: “Sphecidae”) aufgrund morphologischer Merkmale des Exoskeletts. *Zoologische Beiträge* 37: 3–40.
- Panzer, G.W.F. 1801. Nachricht von einem neuen entomologischen Werke, des Hrn. Prof. Jurine in Geneve. *Intelligenzblatt der Literatur-Zeitung [Erlangen]* 1: 160–165. [This work has been suppressed for nomenclatorial purposes: ICZN, 1939: Opinion 135]
- Panzer, G.W.F. 1804a. Fauna Insectorum Germanicae initia, oder Deutschlands Insecten: Heft 85. Nürnberg: Felssecker.
- Panzer, G.W.F. 1804b. Fauna Insectorum Germanicae initia, oder Deutschlands Insecten: Heft 86. Nürnberg: Felssecker.
- Panzer, G.W.F. 1806. Kritische Revision der Insektenfauna Deutschlands [vol. 2]. Nürnberg: Felssecker; [14] + 271 pp., 2 pls.
- Panzer, G.W.F. 1809. Fauna Insectorum Germanicae initia, oder Deutschlands Insecten: Heft 107. Nürnberg: Felssecker.
- Parker, F.D. 1966. A review of the genus *Xenosphex* Williams with biological notes. *Pan-Pacific Entomologist* 42: 190–195.
- Pascoe, F.P. 1870. Contributions towards a knowledge of the Curculionidae. *Journal of the Linnean Society (Zoology)* 10: 434–493.
- Pasteels, J.J. 1969. La systématique générique et subgénérique des Anthidiinae (Hymenoptera, Apoidea, Megachilidae) de l'ancien monde. *Mémoires de la Société Royale d'Entomologique de Belgique* 31: 1–148.
- Pate, V.S.L. 1935. Synonymical notes on the fossorial wasps (Hymenoptera: Sphecidae, Pompilidae and Tiphiidae). *Entomological News* 46: 244–250, 264–267.
- Pate, V.S.L. 1936. *Rhectognathus*, a new group in the *Lindenius* complex (Hymenoptera: Sphecidae: Crabronini). *Entomological News* 47: 147–153.
- Patiny, S. 1999. Etude phylogénétique des Panurginae de l'ancien monde (Hymenoptera, Andrenidae). *Linzer Biologische Beiträge* 31(1): 249–275. [publ. date: 30 July 1999]
- Patton, W.H. 1879. Generic arrangement of the bees allied to *Melissodes* and *Anthophora*. *Bulletin of the United States Geological and Geographical Survey of the Territories* 5: 471–479.

- Perkins, R.C.L. 1908. Some remarkable Australian Hymenoptera. *Proceedings of the Hawaiian Entomological Society* 2: 27–35.
- Perkins, R.C.L. 1912. Notes, with descriptions of new species, on aculeate Hymenoptera of the Australian region. *Annals and Magazine of Natural History*, series 8, 9: 96–121.
- Perty, J.A.M. 1833. [Fascicle 3]. *In* *Delectus Animalium Articulorum, quae in itinere per Brasiliam annis MDCCCXVII–MDCCCXX jussu et auspiciis Maximiliani Josephi I. Bavariae regis augustissimi peracto collegerunt Dr. J. B. de Spix . . . et Dr. C. F. Ph. de Martius. Digessit, descripsit, pingenda curavit Dr. Maximilianus Perty . . . praefactus est et edidit Carol. Frederic. Philip. de Martius . . . Accedit dissertatio de Insectorum in America meridionali habitantium vitae genere, moribus et distributione geographica: 125–224, pls. 25–40. Monachii [Munich]: Hübschmann, [viii] + iii + 44 + 224 pp., 40 pls.*
- Pierce, W.D. 1919. Contributions to our knowledge of the weevils of the superfamily Curculionoidea. *Proceedings of the Entomological Society of Washington* 21(2): 21–38.
- Popov, V.V. 1949. Tribe Pararhopitini (Hymenoptera, Anthophorinae) as an Early Tertiary element of the contemporary fauna of the desert of Central Asia and Egypt. *Doklady Akademii Nauk SSSR* 66(3): 507–510. [In Russian]
- Prentice, M.A. 1998. The comparative morphology and phylogeny of apoid wasps (Hymenoptera: Apoidea) [2 vols.]. Ph.D. diss., University of California, Berkeley, CA.
- Radoszkowski, O. 1868 [1867]. *Materiaux pour servir à l'étude des insectes de la Russie IV. Notes sur quelques Hyménoptères de la tribu des Apides. Horae Societatis Entomologicae Rossicae* 5: 73–90.
- Rasnitsyn, A.P. 1975. Hymenoptera Apocrita of the Mesozoic. *Trudy Paleontologicheskogo Instituta, Akademiya Nauk SSSR* 174: 1–191. [In Russian]
- Rayment, T. 1935. A cluster of bees. Sixty essays on the life-histories of Australian bees, with specific descriptions of over 100 new species, and an introduction by Professor E. F. Phillips. Sydney: Endeavour Press, 752 pp.
- Reed, E.C. 1894. *Entomología [sic] Chilena. Anales de la Universidad [Santiago, Chile]* 85: 599–653.
- Rightmyer, M.G. 2004. Phylogeny and classification of the parasitic bee tribe Epeolini (Hymenoptera: Apidae, Nomadinae). *University of Kansas Museum of Natural History Scientific Papers* 33: 1–49. [publ. date: 6 October 2004]
- Robertson, C. 1901. Some new or little-known bees. *Canadian Entomologist* 33: 229–231.
- Robertson, C. 1902a. Some new or little-known bees, IV. *Canadian Entomologist* 34: 321–331.
- Robertson, C. 1902b. Synopsis of Halictinae. *Canadian Entomologist* 34: 243–250.
- Robertson, C. 1903a. Synopsis of Megachilidae and Bombinae. *Transactions of the American Entomological Society* 29: 163–178.
- Robertson, C. 1903b. Synopsis of Epeolinae. *Canadian Entomologist* 35: 284–288.
- Robertson, C. 1904. Synopsis of Anthophila. *Canadian Entomologist* 36: 37–43.
- Robertson, C. 1922. Synopsis of Panurgidae. *Psyche* 29: 159–173.
- Robertson, C. 1926. Phenology of inquiline and nest-making bees. *Psyche* 33: 116–120.
- Rohwer, S.A. 1916. Sphecoidea. *In* H.L. Viereck, *The Hymenoptera, or wasp-like insects, of Connecticut: 645–697. Mitteilungen der Münchener Entomologischen Gesellschaft* 87: 3–21. [publ. date: 1 December 1997]
- Roig-Alsina, A., and C.D. Michener. 1993. Studies of the phylogeny and classification of long-tongued bees (Hymenoptera: Apoidea). *University of Kansas Science Bulletin* 55(4): 123–162. [publ. date: 30 August 1993]
- de Romand, B. 1840. Sur l'Hyménoptère nommé *Acanthopus Goryi* (voy. cette Revue, 1840, p. 248). *Revue Zoologique par la Société Cuvierienne* 1840: 335–336.
- Rozen, J.G., Jr. 1966. The larvae of the Anthophoridae (Hymenoptera, Apoidea). Part 2. The Nomadinae. *American Museum Novitates* 2244: 1–38. [publ. date: 19 May 1966]
- Rozen, J.G., Jr. 1978. The relationships of the bee subfamily Ctenoplectrinae as revealed by its biology and mature larva (Apoidea: Melittidae). *Journal of the Kansas Entomological Society* 51(4): 637–652. [publ. date: 17 November 1978]
- Rozen, J.G., Jr. 1993. Phylogenetic relationships of *Euherbstia* with other short-tongued bees (Hymenoptera: Apoidea). *American Museum Novitates* 3060: 1–17. [publ. date: 25 March 1993]
- Rozen, J.G., Jr. 1996. First and last larval instars of the cleptoparasitic bee *Hexepeolus rhodogyne* (Hymenoptera: Apidae: Nomadinae). *Memoirs of the Entomological Society of Washington* 17: 188–193.
- Rozen, J.G., Jr. 2003. A new tribe, genus, and species of South American panurgine bee (Andrenidae, Panurginae), oligolectic on *Nolana*

- (Nolanaceae). In G.A.R. Melo and I. Alves-dos-Santos (editors), *Apoidea Neotropica: Homenagem aos 90 Anos de Jesus Santiago Moure*: 93–106. Criciúma: Editora UNESC [Universidade do Extremo Sul Catarinense], xvi + 320 pp.
- Rozen, J.G., Jr., K.R. Eickwort, and G.C. Eickwort. 1978. The bionomics and immature stages of the cleptoparasitic bee genus *Protepeolus* (Anthophoridae, Nomadinae). *American Museum Novitates* 2640: 1–24. [publ. date: 3 January 1978]
- Rozen, J.G., Jr., and R.J. McGinley. 1974. Phylogeny and systematics of Melittidae based on the mature larvae (Insecta, Hymenoptera, Apoidea). *American Museum Novitates* 2545: 1–31. [publ. date: 10 September 1974]
- Rozen, J.G., Jr., and L. Ruz. 1995. South American panurgine bees (Andrenidae: Panurginae), Part II. Adults, immature stages, and biology of *Neffapis longilingua*, a new genus and species with an elongate glossa. *American Museum Novitates* 3136: 1–15. [publ. date 26 May 1995]
- Rozen, J.G., Jr., and D. Yanega. 1999. Nesting biology and immature stages of the South American bee genus *Acamptopoeum* (Hymenoptera: Andrenidae: Panurginae). *University of Kansas Natural History Museum Special Publication* 24: 59–67. [publ. date: 17 December 1999]
- Ruz [-Escudero], L. 1987. Classification and relationships of the panurgine bees (Hymenoptera—Andrenidae). Ph.D. diss., University of Kansas, Lawrence, KS, iii + 312 + [68] pp.
- Ruz, L. 1991. Classification and phylogenetic relationships of the panurgine bees: The Calliopsini and allies (Hymenoptera: Andrenidae). *University of Kansas Science Bulletin* 54(7): 209–256. [publ. date: 31 May 1991]
- Sakagami, S.F. 1974. Sozialstruktur und Polymorphismus bei Furchen- und Schmalbienen. In G.H. Schmidt (editor), *Sozialpolymorphismus bei Insekten: Probleme der Kastenbildung im Tierreich*: 257–293. Stuttgart: Wissenschaftliche Verlagsgesellschaft MBH, xxiv + 974 pp.
- Sakagami, S.F., and C.D. Michener. 1987. Tribes of Xylocopinae and origin of the Apidae (Hymenoptera: Apoidea). *Annals of the Entomological Society of America* 80: 439–450.
- de Saussure, H. 1890 [1892]. Histoire naturelle des Hyménoptères. In A. Grandidier (editor), *Histoire physique, naturelle et politique de Madagascar* [vol. 20]: 1–590. Paris: Imprimerie Nationale, xxi + 590 pp., 27 pls. [NB: date of actual publication is likely 1892]
- Schenck, A. 1859 [1861]. Die nassauischen Bienen. *Revision und Ergänzung der früheren Bearbeitungen. Jahrbücher des Vereins für Naturkunde im Herzogthum Nassau* 14: 1–414. [publ. date: February 1861, *vide* Michener (1986a); also published as Schenck (1861, *vide infra*)]
- Schenck, A. 1860. Verzeichniss der nassauischen Hymenoptera aculeata mit Hinzufügung der übrigen dem Verfasser bekannt gewordenen deutschen Arten. *Stettiner Entomologische Zeitung* 21: 132–157, 417–419.
- Schenck, A. 1861. Die Bienen des Herzogthums Nassau. Wiesbaden: Niedner, iv + 414 + [3] pp. [Publ. date: likely prior to February 1861, *vide* Michener (1986a)]
- Schenck, A. 1866 [1867]. Verzeichniss der nassauischen Hymenoptera aculeata mit Hinzufügung der übrigen deutschen Arten. *Berliner Entomologische Zeitschrift* 10: 317–369. [publ. date: January 1867]
- Schenck, A. 1868 [1869]. Beschreibung der nassauischen Bienen, zweiter Nachtrag. Wiesbaden: Niedner, 1–114 [269–382] pp. [publ. date: before 26 June 1869, *vide* Michener (1986a)]
- Schönherr, C.J. 1836. *Genera et species curculionidum, cum synonymia hujus familiae. Species novae aut hactenus minus cognitae, descriptionibus a Dom. Leonardo Gyllenhal, C. H. Boheman, et entomologis aliis illustratae* [vol. 3, part 2]. Paris: Roret, 506–858 pp.
- Schrottky, C. 1902. Ensaio sobre as abelhas solitárias do Brasil. *Revista do Museo Paulista* 5: 330–613, pls. xii–xiv.
- Schrottky, C. 1909a. Synonymische Bemerkungen über einige südamerikanische Halictinae. *Deutsche Entomologische Zeitschrift* 1909: 479–485.
- Schrottky, C. 1909b. Himenópteros de Catamarca. *Anales de la Sociedad Científica Argentina* 68: 233–272.
- Schrottky, C. 1911. Descrição de abelhas novas do Brasil e der regiões vizinhas. *Revista do Museo Paulista* 8: 71–88.
- Schrottky, C. 1913. La distribución geográfica de los himenópteros argentinos. *Anales de la Sociedad Científica Argentina* 75: 115–144, 180–286.
- Schwarz, M.P., N.J. Bull, and S.J.B. Cooper. 2003. Molecular phylogenetics of allodapine bees, with implications for the evolution of sociality and progressive rearing. *Systematic Biology* 52(1): 1–14.
- Shuckard, W.E. 1840. [Untitled section of Chapter IV: The Hymenoptera]. In W. Swainson and W. E. Shuckard, *On the history and natural arrangement of insects*: 155–190. London: Longman et al., iv + 406 pp.
- Scopoli, J.A. 1770. *Annus Historico Naturalis*.

- Annus IV. Lipsiae [Leipzig]: Hilscher, 150 pp., 2 pls.
- Silveira, F.A. 1995. Phylogenetic relationships and classification of Exomalopsini with a new tribe Teratognathini (Hymenoptera: Apoidea). University of Kansas Science Bulletin 55(12): 425–454. [publ. date: 3 November 1995]
- Silveira, F.A., G.A.R. Melo, and E.A.B. Almeida. 2002. Abelhas Brasileiras: Sistemática e Identificação. Belo Horizonte: Editora IDMAR, 253 pp.
- Smith, F. 1853. Catalogue of the hymenopterous insects in the collection of the British Museum. Part 1. Andrenidae and Apidae. London: British Museum, [i] + 1–198 pp., pls. i–vi.
- Smith, F. 1854. Catalogue of the hymenopterous insects in the collection of the British Museum. Part 2. Apidae. London: British Museum, 199–465 pp., pls. vii–xii.
- Smith, F. 1855. Catalogue of British Hymenoptera in the collection of the British Museum. Part 1. Apidae–bees. London: British Museum, [i] + 252 + 4 pp., pls. a, i–x.
- Smith, F. 1879. Descriptions of new species of Hymenoptera in the collection of the British Museum. London: British Museum, xxi + 240 pp.
- Spinola, M. 1808. Insectorum Liguria species novae aut rariores, quas in agro ligustico nuper detexit, descripsit, et iconibus illustravit Maximilianus Spinola, adjecto catalogo specierum auctoribus jam enumeratarum, quæ in eadem regione passim occurrunt [vol. 2, fasc. 2–4]. Gravier; Genuae [Genoa], ii + 262 pp.
- Spinola, M. 1838 [1839]. Compte-rendu des Hyménoptères, recueilles par M. Fischer pendant son voyage en Egypte. Annales de la Société Entomologique de France 7: 437–546.
- Spinola, M. 1843. Sur quelques Hyménoptères peu connus, recueillis en Espagne, pendant l'année 1842, par M. Victor Ghiliani, voyageur-naturaliste. Annales de la Société Entomologique de France, serie 2 1: 111–144.
- Spinola, M. 1851. Hymenópteros. In C. Gay (editor), Historia física y política de Chile, zoología: tomo sexto [sic] [vol. 6]: 153–569. Paris: Casa del autor, 572 pp.
- Spinola, M. 1853. Compte rendu des hyménoptères inédits provenant du voyage entomologique de M. Ghiliani dans le Para en 1846. Memorie della Reale Accademia delle Scienze di Torino, series 2, 13: 19–94.
- Thomson, C.G. 1869. Opuscula Entomologica [fasc. 1]. Lund: Lundbergska, 82 pp.
- Toro, H. 1980. *Austropanurgus* nuevo género de Panurginae chileno (Hymenoptera, Apoidea). Anales del Museo de Historia Natural de Valparaíso 13: 209–212.
- Toro, H., and A. Moldenke. 1979. Revisión de los Xeromelissinae chilenos (Hymenoptera–Colletidae). Anales del Museo de Historia Natural de Valparaíso 12: 95–182.
- Turner, R.E. 1914. Notes on fossorial Hymenoptera XIII. A revision of the Paranyssoninae. Annals and Magazine of Natural History, series 8, 14: 337–359.
- Turner, R.E. 1915. Notes on fossorial Hymenoptera XV. Annals and Magazine of Natural History, series 8, 15: 62–96.
- Vachal, J. 1905a. *Manuelia*, un nouveau genre d'Hyménoptères mellifères. Bulletin de la Société Entomologique de France 1905: 25–26.
- Vachal, J. 1905b. Un nouveau genre d'Hyménoptères, de la famille Apidae. Bulletin de la Société Entomologique de France 1905: 204.
- Vachal, J. 1909. Espèces nouvelles ou litigieuses d'Apidae du haut bassin du Parana et de régions contiguës et délimitation d'une nouvelle sous-famille Diphaglossinae. Revue d'Entomologie [Caen] 28: 5–72.
- Vachal, J. 1911. Étude sur les *Halictus* d'Amérique (Hym.). Miscellanea Entomologica 19: 9–24, 41–56, 107–112.
- Viereck, H.L. 1916. The Hymenoptera, or wasp-like insects, of Connecticut. Connecticut State Geological and Natural History Survey Bulletin 22: 1–824, pls. i–x.
- Warncke, K. 1977. Ideen zum natürlichen System der Bienen (Hymenoptera, Apoidea). Mitteilungen der Münchener Entomologischen Gesellschaft 67: 39–63.
- Westwood, J.O. 1840. An introduction to the modern classification of insects; found on the natural habits and corresponding organisation of the different families [vol. 2]. London: Longman et al., xi + 587 + 158 pp. [NB: pp. 1–352 appeared in 1839, Melectides appears on 270 and thus dates from 1839]

APPENDIX

HIERARCHICAL CLASSIFICATION OF SUPERFAMILY APOIDEA

Herein I provide a classificatory outline of the superfamily Apoidea. All names, including synonyms (italicized), are included in this listing. Names are provided with their corrected spelling and, therefore, may not appear with the same stem or suffix with which they were originally proposed. Those names for apoid wasps follow Menke (1997) and Menke and Pulawski (2002), with updates from Antropov (2000), but the hierarchical arrangement follows that of Ohl (1996), Prentice (1998), and Melo (1999) so as to reflect phylogenetic relationships relative to the bees. As is frequently done in apoid wasps, I have used subtribes throughout the bees. General references to

the classification of Recent and fossil bees include Michener (2000) and Engel (2001), with excellent accounts of the phylogeny of particular groups by Ascher (2003, 2004), Brady and Danforth (2004), Danforth et al. (2004), Roig-Alsina and Michener (1993), etc. The addition of infratribes or other intercalated ranks (*vide* table 1) could greatly enrich the classificatory hierarchy of bees, more intensely reflecting their presumed phylogenetic relationships. I have not presently felt it warranted to recognize such levels, instead employing only those ranks widely adopted across other insect lineages, particularly elsewhere in the Apoidea.

-
- Superfamily APOIDEA Latreille, 1802a
- “**spheciforms**” [*apoid wasps; paraphyletic grade to Anthophila*]
- Family †ANGAROSPHECIDAE Rasnitsyn, 1975
= †*Baissodidae* Rasnitsyn, 1975
- Family HETEROGYNAIDAE Nagy, 1969, *nomen emendatum* (ICZN, 1987)
- Family AMPULICIDAE Shuckard, 1840
- Subfamily Ampulicinae Shuckard, 1840
- Tribe †Apodolichurini Antropov, 2000
- Tribe †Cretampulicini Antropov, 2000
- Tribe †Mendampulicini Antropov, 2000
- Tribe Dolichurini Lepeletier de Saint Fargeau, 1845
- Tribe Ampulicini Shuckard, 1840
- Family SPHECIDAE Latreille, 1802b
- Subfamily Chloriontinae Fernald, 1905
- Subfamily Sceliphrinae Ashmead, 1899b, *nomen conservandum* (ICZN, 1999a: Art. 40.2)
- Tribe Sceliphriini Ashmead, 1899b, *nomen conservandum* (ICZN, 1999a: Art. 40.2)
= *Pelopoeini* Leach, 1815
- Tribe Podiini de Sausurre, 1892
- Subfamily Stangeellinae Bohart and Menke, 1976
- Subfamily Sphecinae Latreille, 1802b
- Tribe Sphecini Latreille, 1802b
- Tribe Prionychini Bohart and Menke, 1963
- Subfamily Ammophilinae André, 1886
- Family CRABRONIDAE Latreille, 1802b
- Subfamily †Burmastatinae Antropov, 2000 [*Or separate family?*]
- Subfamily Astatinae Lepeletier de Saint Fargeau, 1845
- Tribe Astatini Lepeletier de Saint Fargeau, 1845
= *Diploplectrini* Fox, 1894 [1895]
= *Dimorphini* Brues and Melander, 1932
- Tribe Ammoplanini Evans, 1959
- Tribe Eremiasphecini Menke, 1967
- Subfamily Bembicinae Latreille, 1802b
- Tribe Alyssontini Dalla Torre, 1897
- Tribe Nyssonini Latreille, 1804, *nomen emendatum* (ICZN, 1979)
- Tribe Bembicini Latreille, 1802b
- Subtribe Heliocausina Handlirsch, 1925
- Subtribe Exeirina Dalla Torre, 1897
= *Clitemnestrina* Nemkov and Lelej, 1996
= *Oligina* Nemkov and Lelej, 1996
- Subtribe Gorytina Lepeletier de Saint Fargeau, 1845

- = *Arpactina* Turner, 1915
- = *Hoplisina* Rohwer, 1916
- = *Argogorytina* Nemkov and Lelej, 1996
- Subtribe Handlirschiina Nemkov and Lelej, 1996
- Subtribe Stizina Costa, 1859
- Subtribe Bembicina Latreille, 1802b
- Subtribe Stictiellina Bohart and Horning, 1971
- Subfamily Philanthinae Latreille, 1802b
 - Tribe Philanthini Latreille, 1802b
 - Subtribe Philanthina Latreille, 1802b
 - Subtribe Philanthinina Menke, 1967
 - Tribe Aphilanthopini Bohart, 1966
 - Tribe Pseudoscoliini Menke, 1967
 - Tribe Cercerini Lepeletier de Saint Fargeau, 1845
- Subfamily Pemphredoninae Dahlbom, 1835
 - Tribe Odontosphecini Menke, 1967
 - Tribe Entomosericini Dalla Torre, 1897
 - Tribe Psenini Costa, 1858
 - = *Mimesini* Cresson, 1887
 - = *Psenulini* Gittins, 1969
 - Tribe Pemphredonini Dahlbom, 1835
 - Subtribe Pemphredonina Dahlbom, 1835
 - Subtribe Spilomenina Menke, 1989
 - Subtribe Stigmina Bohart and Menke, 1976
- Subfamily Crabroninae Latreille, 1802b
 - Tribe Mellinini Latreille, 1802b
 - Subtribe Mellinina Latreille, 1802b
 - Subtribe Xenosphecina Parker, 1966
 - Tribe Dinetini Fox, 1894 [1895]
 - Tribe Laphyragogini Bohart and Menke, 1976
 - Tribe Palarini Schrottky, 1909b
 - Tribe Larrini Latreille, 1810
 - Subtribe Larrina Latreille, 1810
 - = *Larradina* de Saussure, 1890 [1892]
 - Subtribe Gastrosericina André, 1886
 - = *Tachytina* Bohart, 1951^a
 - Tribe Miscophini Fox, 1894 [1895]
 - = *Lyrodini* Fox, 1894 [1895]
 - = *Sericophorini* Dalla Torre, 1897
 - = *Nitelini* Dalla Torre, 1897
 - = *Paranyssontini* Turner, 1914
 - Tribe Trypoxylini Lepeletier de Saint Fargeau, 1845
 - = *Pisini* Ashmead, 1899b
 - Tribe Bothynostethini Fox, 1894 [1895]
 - Subtribe Bothynostethina Fox, 1894 [1895]
 - Subtribe Scapheutina Menke, 1968
 - Tribe Oxybelini Leach, 1815
 - Tribe Crabronini Latreille, 1802b
 - Subtribe Anacrabronina Ashmead, 1899b
 - Subtribe Crabronina Latreille, 1802b
 - = *Lindeniina* Ashmead, 1899b
 - = *Thyreopodina* Ashmead, 1899b
 - = *Rhopalina* Ashmead, 1899b
 - = *Soleniina* Bradley, 1926
 - = *Pemphilidina* Pate, 1935
 - = *Karossiina* Pate, 1936

Family Incertae Sedis

- Subfamily †Cirrosphecinae Antropov, 2000 (*perhaps near Ampulicidae?*)

Anthophila Latreille, 1804 *vel* 1807 [*bees*]^b

- Family COLLETIDAE Lepeletier de Saint Fargeau, 1841

- Subfamily Stenotritinae Cockerell, 1934
- Subfamily Diphaglossinae Vachal, 1909
 - Supertribe Caupolicaniti Michener, 1944
 - Tribe Caupolicanini Michener, 1944
 - Supertribe Diphaglossiti Vachal, 1909
 - Tribe Diphaglossini Vachal, 1909
 - Tribe Dissoglottini Moure, 1945
 - = *Ptiloglossidiini* Moure, 1953
 - = *Mydrosomatini* Michener, 1966
- Subfamily Paracolletinae Cockerell, 1934, *nomen protectum* (ICZN, 1993)
 - = *Neopasiphaeinae* Cockerell, 1930
 - = *Phenacolletinae* Cockerell, 1934, *nomen nudum*
 - = *Eulonchopriinae* Moure, 1945
 - = *Lonchopriinae* Moure, 1945
- Subfamily Colletinae Lepeletier de Saint Fargeau, 1841
- Subfamily Scraptrinae Ascher and Engel, *subfamilia novum*
- Subfamily Hylaeinae Viereck, 1916, *nomen protectum* (ICZN, 1993)
 - Supertribe Hylaeiti Viereck, 1916, *nomen protectum* (ICZN, 1993)
 - Tribe Euryglossini Michener, 1944
 - Tribe Hylaeini Viereck, 1916, *nomen protectum* (ICZN, 1993)
 - = *Prosopidini* Fallén, 1813
 - = *Palaeorhizini* Perkins, 1908
 - = *Meroglossini* Perkins, 1908
 - = *Hyleoidini* Cockerell, 1930a
 - Supertribe Xeromelissiti Cockerell, 1926
 - Tribe Xeromelissini Cockerell, 1926
 - = *Chilicolini* Michener, 1944
- Family HALICTIDAE Thomson, 1869, *nomen protectum* (ICZN, 1993)
 - Subfamily Rophitinae Schenck, 1866 [1867]
 - Tribe Rophitini Schenck, 1866 [1867]
 - = *Dufoureini* Robertson, 1904
 - = *Halictoidini* Börner, 1919
 - = *Systrophini* Handlirsch, 1925
 - Tribe Penapini Engel, 2001
 - Subfamily Nomiinae Robertson, 1904
 - Subfamily Halictinae Thomson, 1869, *nomen protectum* (ICZN, 1993)
 - Supertribe Nomioiditi Börner, 1919
 - Tribe Nomioidini Börner, 1919
 - Supertribe Halictiti Thomson, 1869, *nomen protectum* (ICZN, 1993)
 - Tribe Halictini Thomson, 1869, *nomen protectum* (ICZN, 1993)
 - Subtribe Sphecodina Schenck, 1868 [1869]
 - Subtribe Halictina Thomson, 1869
 - = *Gastrohalictina* Schrottky, 1911
 - = *Lasioglossina* Robertson, 1926
 - = *Chloralictina* Robertson, 1926
 - = *Thrinchostomatina* Sakagami, 1974
 - Tribe Caenohalictini Michener, 1954
 - Subtribe Agapostemonina Engel, 2000
 - Subtribe Caenohalictina Michener, 1954
 - Tribe Augochlorini Beebe, 1925, *nomen protectum* (ICZN, 2000)
 - Subtribe Corynurina Engel, 2000
 - Subtribe Augochlorina Beebe, 1925, *nomen protectum* (ICZN, 2000)
 - = *Oxystoglossina* Schrottky, 1909a
 - = *Augochloropsina* Moure, 1943
 - = *Megaloptina* Moure, 1943
 - = *Ischnomelissina* Engel, 2000
 - = *Thectochlorina* Engel, 2000
 - = *Megaloptidiina* Engel, 2000
- Family ANDRENIDAE Latreille, 1802a
 - Subfamily Euherbstiinae Moure, 1950

- Subfamily Andreninae Latreille, 1802a
 - = *Anthreninae* Burmeister, 1829
 - = *Alocandreninae* Michener, 2000
- Subfamily Oxaeinae Ashmead, 1899a
- Subfamily Panurginae Leach, 1815
 - Supertribe Nolanomelissiti Rozen and Ascher *In* Rozen, 2003
 - Tribe Nolanomelissini Rozen and Ascher *In* Rozen, 2003
 - Supertribe Panurgiti Leach, 1815
 - Tribe Calliopsini Robertson, 1922
 - Tribe Protandrenini Robertson, 1904
 - = *Liphanthini* Michener, 2000, *nomen nudum*
 - = *Austropanurgini* Michener, 2000, *nomen nudum*
 - Tribe Neffapini Ascher, *tribus novum*
 - Tribe Protomeliturgini Ruz, 1991
 - Tribe Meliturgulini Engel, 2001
 - Tribe Panurgini Leach, 1815
 - Subtribe Melitturgina Newman, 1834
 - = *Mermiglossina* Patiny, 1999
 - Subtribe Camptopoeina Patiny, 1999
 - Subtribe Panurginina Patiny, 1999
 - Subtribe Panurgina Leach, 1815
 - Subtribe Perditina Robertson, 1922
- Family †PALEOMELITTIDAE Engel, 2001
- Family MELITTIDAE Schenck, 1860
 - Subfamily Meganomiinae Michener, 1981
 - Subfamily Macropidinae Robertson, 1904
 - Tribe †Eomacropidini Engel, 2001
 - Tribe Macropidini Robertson, 1904
 - Subfamily Melittinae Schenck, 1860
 - Tribe Redivivini Engel, 2001
 - Tribe Melittini Schenck, 1860
 - Subfamily Dasypodainae Börner, 1919
 - Supertribe Promelittiti Michener, 1981
 - Tribe Promelittini Michener, 1981
 - Supertribe Dasypodaiti Börner, 1919
 - Tribe Afrodasypodini Engel, *tribus novum*
 - Tribe Sambini Michener, 1981
 - Tribe Dasypodaini Börner, 1919
 - Subtribe Dasypodaina Börner, 1919
 - Subtribe Hesperapina Ascher and Engel, *subtribus novum*
- Family MEGACHILIDAE Latreille, 1802b
 - Subfamily Pararhophitinae Popov, 1949
 - Subfamily Fideliinae Cockerell, 1932
 - Tribe Neofideliini Engel, 2004
 - Tribe Fideliini Cockerell, 1932
 - Subfamily Lithurginae Newman, 1834
 - Tribe †Protolithurgini Engel, 2001
 - Tribe Lithurgini Newman, 1834
 - = *Trichothurgini* Moure, 1949
 - Subfamily Megachilinae Latreille, 1802b
 - Tribe Anthidiini Ashmead, 1899a, *nomen protectum* (ICZN, 1993)
 - Subtribe Anthidiina Ashmead, 1899a, *nomen protectum* (ICZN, 1993)
 - = *Stelididina* Schenck, 1860
 - = *Trachusina* Robertson, 1904
 - = *Dianthidiina* Moure, 1947
 - = *Pachyanthidiina* Pasteels, 1969
 - = *Icteranthidiina* Pasteels, 1969
 - = *Euaspinga* Pasteels, 1969
 - Subtribe Dioxyina Cockerell, 1902
 - Tribe †Glyptapini Cockerell, 1909b

- Tribe †Ctenoplectrellini Engel, 2001
- Tribe Osmiini Newman, 1834
 - Subtribe Heriadinina Michener, 1941
 - = *Trypetina Robertson, 1903a, nomen praeoccupatum*
 - Subtribe Osmiina Newman, 1834
 - = *Chelostomina Kirby In Kirby and Spence, 1837*
 - = *Hoplitina Nobile and Tomarchio, 2000, nomen nudum*
- Tribe Megachilini Latreille, 1802b
 - = *Coelioxyni Dalla Torre and Fries, 1894*
 - = †*Chalicodomopsini Engel, 1999b, nomen nudum*
- Family APIDAE Latreille, 1802a
 - Subfamily Xylocopinae Latreille, 1802b
 - Supertribe Xylocopiti Latreille, 1802b
 - Tribe Xylocopini Latreille, 1802b
 - Supertribe Manuelliiti Sakagami and Michener, 1987
 - Tribe Manuelliini Sakagami and Michener, 1987
 - Supertribe Ceratiniti Latreille, 1802b
 - Tribe Ceratinini Latreille, 1802b
 - Tribe †Boreallodapini Engel, 2001
 - Tribe Allodapini Cockerell, 1902
 - Subtribe Macrogaleina Engel, *subtribus novum*
 - Subtribe Allodapina Cockerell, 1902
 - = *Exoneurina Robertson, 1904*
 - = *Eucondylopsini Friese, 1909*
 - Subfamily Nomadinae Latreille, 1802a
 - Supertribe Neolarriti Fox, 1894 [1895]
 - Tribe Neolarrini Fox, 1894 [1895]
 - Tribe Ammobatini Handlirsch, 1925, *nomen protectum* (ICZN, 1993)
 - Subtribe Pasitina Robertson, 1926
 - Subtribe Ammobatina Handlirsch, 1925, *nomen protectum* (ICZN, 1993)
 - = *Phileremina Lepeletier de Saint Fargeau, 1841*
 - = *Melanempidina Handlirsch, 1925*
 - Tribe Caenoprosopidini Michener, 1944
 - Supertribe Nomaditi Latreille, 1802a
 - Tribe Townsendiellini Michener, 1944
 - Tribe Nomadini Latreille, 1802a
 - Tribe Biastini Linsley and Michener, 1939
 - = *Neopasitini Linsley and Michener, 1939*
 - Tribe Ammobatoidini Michener, 1944
 - = *Holcopasitini Rozen, 1966*
 - Tribe Hexepeolini Rozen, 1996
 - Tribe Brachynomadini Roig-Alsina and Michener, 1993
 - Tribe Epeolini Robertson, 1903b
 - Subtribe Odyneropsina Handlirsch, 1925
 - Subtribe Rhogepeolina Rightmyer, 2004
 - Subtribe Epeolina Robertson, 1903b
 - Subtribe Thalestriina Rightmyer, 2004
 - Subfamily Apinae Latreille, 1802a
 - Supertribe Euceriti Latreille, 1802b
 - Tribe Osirini Handlirsch, 1925 [*allied to Tapinotaspidini?; supertribal placement tentative*]
 - = *Epeoloidini Linsley and Michener, 1939*
 - Tribe Isepeolini Rozen, Eickwort, and Eickwort, 1978
 - Tribe Protepeolini Linsley and Michener, 1939
 - Tribe Exomalopsini Vachal, 1909
 - Tribe Ancylini Michener, 1944
 - Tribe Teratognathini Silveira, 1995
 - Tribe Eucerini Latreille, 1802b
 - Subtribe Eucerinodina Michener and Moure, 1957
 - Subtribe Eucerina Latreille, 1802b
 - = *Melissodina Robertson, 1901*

- = *Tetraloniina* Schrottky, 1913
- = *Canephorulina* Michener, LaBerge, and Moure, 1955
- Tribe Ctenoplectrini Cockerell, 1930a
- Tribe Emphorini Robertson, 1904
 - Subtribe Ancyloscelidina Engel and Michener, *subtribus novum*
 - Subtribe Emphorina Robertson, 1904
 - = *Melitomina* Vachal, 1909
 - = *Entechniina* Cockerell, 1906
- Tribe Tapinotaspidini Moure, 1992 [1994]
 - = *Paratetrapediini* Moure, 1992 [1994]
- Supertribe Apiti Latreille, 1802a
 - Tribe Tetrapediini Michener and Moure, 1957 [*supertribal placement tentative*]
 - Tribe Rhathymini Lepeletier de Saint Fargeau, 1841
 - Tribe Anthophorini Dahlbom, 1835
 - = *Podaliriini* Latreille, 1802b
 - = *Megillini* Thomson, 1869
 - = *Habropodini* Marikovskaya, 1976
 - Tribe Melectini Westwood, 1840 [1839]
 - Tribe Ericrocidini Cockerell and Atkins, 1902
 - = *Ctenioschelini* Michener, 1965a, *nomen nudum*
 - Tribe Centridini Cockerell and Cockerell, 1901
 - = *Hemisiini* Cockerell and Robbins, 1910
 - = *Epicharitini* Schrottky, 1913
 - Corbiculate apines* [*Corbiculata* Engel, 1998b^c]—
 - Tribe Euglossini Latreille, 1802b
 - = *Eulaemini* Lepeletier de Saint Fargeau, 1841
 - = *Exaeretini* Cockerell, 1908a
 - = *Chrysanthedini* Handlirsch, 1925
 - = *Aglaini* Handlirsch, 1925
 - = *Eufrieseini* Moure, 1944
 - Tribe Bombini Latreille, 1802b
 - = *Apathini* Newman, 1834
 - = *Psithyrini* Lepeletier de Saint Fargeau, 1841
 - = *Bremeni* Frison, 1919
 - Tribe †Electrobombini Engel, 2001
 - Tribe †Electrapini Engel, 1998a
 - Tribe Apini Latreille, 1802a
 - Tribe †Melikertini Engel, 2001
 - Tribe Meliponini Lepeletier de Saint Fargeau, 1836
 - = *Trigonini* Moure, 1946
 - = *Lestrimelittini* Moure, 1946

† Fossil taxon.

^a The availability of this name from Bohart (1951) should, perhaps, be reinvestigated as the few statements on p. 945 where the name is proposed can only marginally be considered to constitute a differential diagnosis. More likely, the name was truly made available in later papers. Given that Tachytini is a synonym it is merely an academic point.

^b The name Anthophila may be dated from either its proposal as a plural (Latreille, 1804) or its later emendation (Latreille, 1807).

^c I created the nonranked name “Corbiculata” for a 1998 symposium on corbiculate bee relationships (Engel, 1998b). Although it was picked up by some of the symposium attendants and organizers, I do not recommend its adoption and instead favor the informal terminology of simply “corbiculate bees” for this group.

Complete lists of all issues of the *Novitates* and the *Bulletin* are available at World Wide Web site <http://library.amnh.org/pubs>. Inquire about ordering printed copies via e-mail from scipubs@amnh.org or via standard mail from: American Museum of Natural History, Library—Scientific Publications, Central Park West at 79th St., New York, NY 10024. TEL: (212) 769-5545. FAX: (212) 769-5009.